

Are we doing alright?

Realities of violence, mental health and access to healthcare related to sexual orientation and gender identity and expression in eSwatini

RESEARCH REPORT BASED ON A COMMUNITY-LED STUDY IN NINE AFRICAN COUNTRIES

ALEX MÜLLER, KRISTEN DASKILEWICZ AND THE SOUTHERN AND EAST AFRICAN RESEARCH COLLECTIVE ON HEALTH (SEARCH)



To cite this report:

Müller, A., Daskilewicz, K. and the Southern and East African Research Collective on Health (2019). 'Are we doing alright? Realities of violence, mental health, and access to healthcare related to sexual orientation and gender identity and expression in eSwatini: Research report based on a community-led study in nine countries'. Amsterdam: COC Netherlands.

Contributors to the eSwatini country report:

The research team in eSwatini consisted of Thuthu Magagula, Sindy Matse, Melusi Simelane and Sanele Sibiya, as well as fieldworkers from The Rock of Hope.

This report is part of a series of ten reports.




This publication is published under Creative Commons 4.0

You are free to share, copy and redistribute the material in any medium or format under the following conditions:

- Attribution** — You must attribute the work in the manner specified above (but not in any way that suggests the author endorses you or your use of the work).
- Non-commercial** — You may not use the material for commercial purposes.
- No derivatives** — If you remix, transform, or build upon the material, you may not distribute the modified material.

This document is published jointly by COC Netherlands and the Southern and East African partners of COC, who receive funding from the Ministry of Foreign Affairs of The Netherlands.



Are we doing alright?

Realities of violence, mental health and access to healthcare related to sexual orientation and gender identity and expression in eSwatini

RESEARCH REPORT BASED ON A COMMUNITY-LED STUDY IN NINE AFRICAN COUNTRIES

ALEX MÜLLER, KRISTEN DASKILEWICZ AND THE SOUTHERN AND EAST AFRICAN RESEARCH COLLECTIVE ON HEALTH (SEARCH)



The SEARCH Collective

Zimbabwe



Gays and Lesbians of Zimbabwe



Sexual Rights Centre

Botswana



Bonela



Lesbians, Gays and Bisexuals of Botswana



Rainbow Identity Association

Zambia



Friends of Rainka



The Lotus Identity



TransBantu Zambia

Netherlands



COC



South Africa



Durban Lesbian and Gay Community and Health Centre



Gender Dynamix



Gender Health and Justice Research Unit, University of Cape Town



OUT LGBT Well-Being



Triangle Project

Lesotho

The People's Matrix Association



Ethiopia

Two organisations (names withheld)

Kenya

Ishtar-MSM



Jinsiangu



Maaygo



Minority Womyn in Action



National Gay and Lesbian Human Rights Commission



People Marginalised and Aggrieved



eSwatini

The Rock of Hope



Malawi

Centre for the Development of People



ACKNOWLEDGMENTS

To conceptualise, coordinate, implement, analyse, write and disseminate a large, multi-site study through collaboration and partnership would not have been possible without the immense support and dedication of many, many people, within and outside of the SEARCH collective.

First and foremost, we thank all research participants, who shared their time and were willing to answer difficult questions, even when there were potential risks involved at times.

The research coordinators, research assistants and fieldworkers from all civil society organisations involved worked tirelessly to find and interview participants. Thank you: without you this project would not have been possible.

Our Study Oversight Committee provided guidance and on-going opportunity for reflection. Thank you to Prof Lillian Artz, Jillian Butterworth, Lucinda van der Heever, Bram Langen, Dr Ingrid Lynch, Dr Olusegun Odumosu and Liesl Theron.

We would like to thank Dr Anneli Hardy, who has provided expertise and support for the statistical analysis; our colleagues Dr Aisling Heath, Talia Meer, Vuyelwa Mehlomakhulu and Dr Virginia Zweigenthal, who read and commented on earlier versions of this report; and our research interns and assistants, Mina Haji, Marie Lunau, Lilian Mubangizi, Agatha Nyambi and Leslene Stott, who assisted with countless small and not-so-small tasks that enabled the work daily.

We especially thank our colleagues who supported this project as local co-investigators: Sindy Matse from the National AIDS Council in eSwatini, Dr Chelsea Morrioni from the Botswana UPenn Partnership and the Liverpool School of Tropical Medicine, Prof Adamson Muula from the College of Medicine at the University of Malawi, and Nelson Muparamoto from the Department of Sociology at the University of Zimbabwe.

There have been many others who have shared insights, and given practical, intellectual and emotional support through this long, sometimes difficult, but fruitful process, and we are grateful to everyone who has supported this project over the last several years across countries and continents.

This work has truly been the product of queer labour, and whilst the report documents the manifold challenges faced by LGBTI people in East and Southern Africa, it is equally testament to our mutual care, our resilience, resourcefulness and agency.

Contents

The SEARCH Collective	2
Acknowledgments	4
REPORT SUMMARY	10
Introductory comments	10
Sexual orientation, gender identity and expression and minority stress	12
Sexual orientation, gender identity and expression and structural stigma	13
The structure of this report.....	14
METHODOLOGY	15
Participatory approach	15
Study design.....	17
Design of study aims	17
The survey	17
Fieldworker training	20
Who could participate in the survey?.....	21
Sampling methodology	21
Collecting data	22
Pilot study	22
Analysing data	23
Research approvals and regulatory compliance.....	23
FINDINGS IN ESWATINI.....	24
Sexual orientation, gender identity and expression in eSwatini	24
The study population: sample characteristics	25
Sociodemographic characteristics	25
Sexual and gender diversity / sexual orientation and gender identity	26
Socioeconomic circumstances	29
Social support and being 'out'	31
Discrimination in healthcare.....	32
Health-seeking behaviour	34
Gender affirming care	35
Experiences of violence.....	36
Verbal harassment	37
Sexual violence	37
Physical violence	40
Perpetrators of violence.....	43
Impact of violence	45
Reporting of violence	45
Mental health outcomes.....	46
Mental health outcomes in the overall sample	46
Depression	48
Anxiety.....	51

Suicidality	53
Alcohol use.....	55
Drug use	58
Tobacco use	60
Experiences of violence and health outcomes of lesbian participants	62
Experiences of violence and health outcomes of gay participants	65
Experiences of violence and health outcomes of bisexual participants	68
Experiences of violence and health outcomes of gender minority participants	71
Gender affirming care	71
Health outcomes	72
LIMITATIONS	75
CONCLUSION	76
RECOMMENDATIONS	77
Recommendations for national government.....	77
Recommendations for civil society organisations	77
Recommendations for donors	78
Recommendations for academics and researchers.....	78
REFERENCES	79
GLOSSARY OF TERMS RELATED TO SEXUAL ORIENTATION, GENDER IDENTITY AND EXPRESSION	83
GLOSSARY OF TERMS RELATED TO THE STATISTICAL ANALYSIS	85
APPENDIX 1: DETAILED METHODOLOGY	87
Measures: Sexual orientation and gender identity	87
Survey questions.....	87
Categorisation for analysis.....	87
Measures: Mental health	89
CES-D 10: Depression.....	89
GAD-7: Anxiety	89
AUDIT: Alcohol.....	89
DUDIT: Drugs	90
Signs of post-traumatic stress.....	90
Sampling and enrolment.....	90
Data management	93
Data quality	93
Data cleaning.....	93
Conflicting data	94
Data analysis.....	94
Describing the data	94
Measuring associations	94
Logistic regression.....	95
Missing data.....	96
APPENDIX 2: QUESTIONNAIRE	98

List of tables

- Table 1: Community partner organisations..... 16
- Table 2: Number of participants, by country22
- Table 3: Research approvals.....23
- Table 4: Sociodemographic characteristics26
- Table 5: Participant sexual orientation and gender identity, self-identified27
- Table 6: Social and financial capital.....29
- Table 7: Social support and being 'out'31
- Table 8: Healthcare access and discrimination.....32
- Table 9: Gender affirming practices35
- Table 10: Access to gender affirming care.....36
- Table 11: Experiences of harassment and violence36
- Table 12: Perpetrators of lifetime sexual and physical violence44
- Table 13: Reporting violence45
- Table 14: Overall mental health outcomes46
- Table 15: Health outcomes and experiences of violence of lesbian participants62
- Table 16: Health outcomes and experiences of violence of gay participants65
- Table 17: Health outcomes and experiences of violence of bisexual participants.....68
- Table 18: Gender affirming practices71
- Table 19: Health outcomes and experiences of violence of
gender minority participants72

List of Figures

Figure 1:	eSwatini participant sample.....	25
Figure 3:	Participants' gender identities.....	29
Figure 4:	Health-seeking behaviour in previous year	34
Figure 5:	Verbal harassment, past year.....	37
Figure 6:	Sexual violence, lifetime.....	38
Figure 7:	Sexual violence, past year	38
Figure 8:	Sexual violence, by sexual orientation.....	39
Figure 9:	Sexual violence, by gender identity.....	39
Figure 10:	Physical violence, lifetime	41
Figure 11:	Physical violence, past year.....	41
Figure 12:	Physical violence, by sexual orientation.....	42
Figure 13:	Physical violence, by gender identity.....	43
Figure 14:	Depression in overall sample.....	48
Figure 15:	Depression diagnosis and treatment.....	49
Figure 16:	Depression, by sexual orientation	50
Figure 17:	Depression, by gender identity	50
Figure 18:	Signs of anxiety, overall sample.....	51
Figure 19:	Anxiety diagnosis and treatment.....	51
Figure 20:	Anxiety levels, by sexual orientation.....	52
Figure 21:	Anxiety levels, by gender identity	53
Figure 22:	Suicidal ideation, overall sample.....	53
Figure 23:	Suicide attempts, lifetime and past year	54
Figure 24:	Suicide attempts, by sexual orientation.....	54
Figure 25:	Suicide attempts, by gender identity.....	55
Figure 26:	Alcohol use, overall sample	56
Figure 27:	Alcohol use, by sexual orientation	56
Figure 28:	Alcohol use, by gender identity	57
Figure 29:	Drugs use, overall sample.....	58
Figure 30:	Drug use, by sexual orientation.....	59
Figure 31:	Drug use, by gender identity.....	59
Figure 32:	Tobacco use, overall sample.....	60
Figure 33:	Tobacco use, by sexual orientation.....	61
Figure 34:	Tobacco use, by gender identity	61
Figure 35:	Gender identities of lesbian participants	62
Figure 36:	Gender identities of gay participants	65
Figure 37:	Gender identities of bisexual participants.....	68
Figure 38:	Sexual orientations of gender minority participants.....	71

LIST OF ACRONYMS

ACHPR	African Commission for Human and People's Rights
AOR	Adjusted odds ratio
AUDIT	Alcohol Use Disorders Identification Test
CBPR	Community-based participatory research
CEDEP	Centre for the Development of People
CES-D10	10-item Centre for Epidemiological Studies Depression Scale
CI	Confidence interval
COC	Cultuur en Ontspanningscentrum (Center for Culture and Leisure)
DSM	Diagnostic and statistical manual of mental disorders
DUDIT	Drug Use Disorders Identification Test
EDMS	Electronic Data Management System
GAD-7	Generalized Anxiety Disorder 7-item scale
GALZ	Gays and Lesbians of Zimbabwe
GATE	Global Action for Trans* Equality
GHJRU	Gender Health and Justice Research Unit
GNC	Gender non-conforming
HCT/ HIV VCT	HIV voluntary testing and counselling
ICD	International Classification of Disease
LGBT	Lesbian, Gay, Bisexual and Transgender
LGBTI	Lesbian, Gay, Bisexual and Transgender and Intersex
MSM	Men who have sex with men
NGLHRC	National Gay and Lesbian Human Rights Commission
NGO	Non-governmental organisation
n	Sample size
p	p value
SGM	Sexual and gender minority
SOGI	Sexual orientation and gender identity
SOGIE	Sexual orientation and gender identity and expression
SRC	Sexual Rights Centre
STI	Sexually transmitted infection
TBZ	Trans Bantu Zambia
UCT	University of Cape Town
US	United States
WHO	World Health Organization
WSW	Women who have sex with women

REPORT SUMMARY

This report presents research findings on the mental health and well-being of lesbian, gay, bisexual, transgender and intersex (LGBTI) people in eSwatini. It also presents findings on LGBTI people's experiences of violence, and experiences in accessing healthcare.

It is part of a series of reports based on research in nine countries of Southern and East Africa: in Botswana, Ethiopia, Kenya, Lesotho, Malawi, South Africa, eSwatini, Zambia and Zimbabwe. The research was done collaboratively by a consortium of non-governmental organisations (NGOs), academic researchers from the University of Cape Town, and COC Netherlands who funded the project and provided logistical support.

Across those nine countries, we used a standardised questionnaire to survey 3,796 people, and ask about physical and sexual violence, depression, anxiety, suicidality and substance use, as well as experiences of discrimination when accessing healthcare.

The findings give us a sense of the precarious state of LGBTI people's mental health and well-being in East and Southern Africa, and the high levels of violence that LGBTI people experience: compared to what we know from the general population, LGBTI people have higher levels of mental health concerns, have experienced more violence, and have faced barriers to healthcare that are directly linked to their sexual orientation, gender identity or gender expression.

Our findings show that in the nine countries of this study, as elsewhere in the world, discrimination, stigma and marginalisation related to sexual orientation, gender identity and gender expression place LGBTI people at higher risk for mental health concerns and violence.

Introductory comments

Over the last two decades research on lesbian, gay, bisexual and transgender persons, health and violence has highlighted substantial vulnerabilities and health disparities based on sexual orientation, and gender identity and expression in many parts of the world. There is growing awareness of the broad ranging negative consequences of stigma, marginalization and discrimination on the health of people who identify as, or are perceived to be, lesbian, gay, bisexual, transgender and gender diverse (LGBT) (Mayer *et al.*, 2008; Institute of Medicine, 2011; Logie, 2012; Pega and Veale, 2015). For example, in a recent landmark report on LGBT health (Institute of Medicine, 2011), the United States Institute of Medicine pointed out that LGBT people are at increased risk of violence, harassment, and victimization. These findings underscore the link between stigma, marginalization and discrimination and corroborate that sexual orientation, gender identity and expression are important determinants of vulnerability and health (Logie, 2012; Pega and Veale, 2015).

LGBT people are not a homogenous population. The acronyms LGBT or LGBTI (“I” for intersex¹) group individuals together based on similar experiences of discriminatory treatment in society because they fall outside of social norms about sexuality and gender, due to their sexual orientation, gender identity, gender expression, and/or sex characteristics. While this is helpful to analyse the consequences of marginalization, it is important not to assume that individuals under this umbrella acronym necessarily have similar experiences or needs. In fact, individual experiences differ greatly across the populations covered under the acronym. Thus, the populations represented by each individual letter in the acronym are complex and heterogeneous, even more so when differences in race, age, ability, religion, culture, socioeconomic class, and geographic location are also taken into account. In this report, we use the acronym LGBTI in order to point to similar experiences of stigma, marginalization and discrimination based on sexual orientation, gender identity, gender expression and sex characteristics in heteronormative societal frameworks. However, frequently we disaggregate this umbrella into its constituent groups in order to highlight specific characteristics and differences.

Until 1973, the American Psychological Association considered same-sex orientation, attraction, and behaviour (formerly referred to narrowly as homosexuality) to be a mental illness. It is now widely recognised that what is considered a mental illness depends on what society and scientists at a certain time and in a certain context agree to be ‘abnormal’ behaviours, cognitions and emotions (Gergen, 2001). Today, international medical and health organisations, such as the World Psychiatry Association have clearly stated that same-sex orientation, attraction, and behaviour are not mental illnesses, and that attempts to ‘treat’ same-sex sexual orientation are harmful and without evidence of success (Bhugra *et al.*, 2016). The South African Society of Psychiatrists agrees that “there is no scientific evidence that reparative or conversion therapy is effective in changing a person’s sexual orientation. There is, however, evidence that this type of therapy can be destructive” (Victor *et al.*, 2014). Further, in 2015 a panel of experts from the Academy of Science of South Africa, endorsed by the Uganda National Academy of Sciences, condemned the use of ‘conversion’ therapy and called for widespread interventions to generate support for LGBTI people, particularly among healthcare providers (Academy of Science of South Africa, 2015).

Gender variance or diversity (formerly called non-conforming or transgender gender identity), unlike same-sex sexual orientation, remains classified as a mental illness by the American Psychological Association. Many argue that this is for the same reasons that same-sex sexual orientation was once classified as a mental illness (Drescher, 2015), and that gender variance is not pathological (Kara, 2017; Suess Schwend *et al.*, 2018). In the process of revising the International Classification of Disease (ICD), the World Health Organisation is thus proposing to remove the diagnosis related to gender variance from the list of mental health conditions (De Cuypere and Winter, 2016; Robles *et al.*, 2016; World Health Organization, 2018a).

1 People with diverse sex characteristics, (also referred to as ‘intersex’) share similar experiences of discrimination and marginalisation as people with non-normative sexual orientations, gender identities and expressions. Additionally, people with diverse sex characteristics often have experienced forced genital mutilation by healthcare providers, and experience the physical, psychological and emotional consequences thereof. It was outside the scope of this research project to investigate these forced treatments. We strongly recommend that specific research into forced genital mutilations, and the impact of those on people with diverse sex characteristics, be done.

Diversity in sex characteristics (formerly called 'intersex'), like gender variance, remains classified as a pathological condition in the current classification of disease (World Health Organization, 2018b). Like for gender variance, many argue that this is a reflection of social attitudes towards diversity in sex characteristics, that such diversity is not per se pathological, and that regarding diversity of sex characteristics as a pathology increases the vulnerability of people to forced genital surgery, which is recognised as unlawful (GATE, 2017).

Sexual orientation, gender identity and expression and minority stress

Now that it is widely understood that same-sex sexual orientation and gender variance are not mental illnesses themselves, researchers have started to look at the mental health and well-being of people who identify as lesbian, gay, bisexual, transgender and intersex. Whilst this work is largely based in the US, the circumstances of minority stress for people on the African continent may not be all that different, and it is useful to know about the work that has already been done in the US in order to contextualise and interpret the findings of this report.

Researchers have found that compared with their heterosexual, cisgender counterparts, sexual and gender minority² populations suffer from more mental health problems, such as substance use (including alcohol, tobacco and illegal drug use), affective disorders (for example, depression and anxiety disorders) and suicide (Meyer, 2003; Hendricks and Testa, 2012; Bockting *et al.*, 2013a). The reason for these disparities in mental health outcomes is that stigma (widespread disapproval held by many people in a society), prejudice, discrimination and structural stigma (social stigma that is institutionalised or made into law, such as laws that criminalise consensual same-sex behaviour), lead to stressful social environments for sexual and gender minorities (Meyer, 2003; Hendricks and Testa, 2012; Hatzenbuehler *et al.*, 2014). This is called minority stress.

Meyer (2003) points out that minority stress adds to general stress that all people experience. It is chronic – that is it lasts a long time, or a person's entire life, as it is linked to underlying social and cultural norms (and stigma) that are relatively stable and only change slowly, if at all. Lastly, minority stress is socially based – that means it stems from social processes, institutions and structures (for example, laws that criminalise consensual same-sex activity), and not from individual events (such as change in financial circumstances, or death of a loved one).

Meyer (2003) also explains how minority stress affects people with same-sex sexual orientation, attraction, and behaviour, and suggests that there are four different processes that contribute to minority stress and mental health problems among sexual minorities. First, chronic and acute events or social circumstances might add to stress. This might include experiences of discrimination in healthcare facilities or schools, or being insulted or harassed in private or public. Second, expecting such stressful events, and guarding oneself against them, also leads to stress (regardless of whether or not the discriminatory encounter actually happens). Third, hearing negative, discriminatory attitudes means that people internalise the idea that they have less value. And fourth, hiding one's sexual orientation in anticipation of discriminatory events further contributes to stress.

2 For the purposes of this report, gender minority people are those who do not identify as cisgender, and are inclusive of the following: those who self-identify as transgender, gender non-conforming (GNC) or non-binary, have a different gender identity from what was assigned to them at birth, and/or identify as intersex.

Hendricks and Testa (2012) explain how minority stress affects gender minority people, and argue that the same factors shape minority stress for this group. That is, as with same-sex sexual orientation, it is not gender variance itself that is a mental illness, but that, essentially, “hostile and stressful social environments” (p. 462) lead to an increase in mental health problems among gender minority people.

Sexual orientation, gender identity and expression and structural stigma

Stigma against same-sex orientation and gender variance is one of the key factors that underlie the stressors in the minority stress model. A recent study built on the work by Meyer (2003) and Hendricks and Testa (2012) and examined the impact stigma has on the health and well-being of sexual minority³ people. This study specifically looked at the impact of structural stigma, defined as social prejudice against lesbian, bisexual and gay people at the community level. This study found that sexual minorities who lived in areas with high structural stigma in the United States were three times more likely to die from homicide and violence-related deaths, when compared to sexual minority people living in areas with low structural stigma (Hatzenbuehler *et al.*, 2014), though this was later shown not to be statistically significant (Hatzenbuehler *et al.*, 2018). The study also showed that sexual minorities in high-stigma areas were more likely to die from suicide. Additionally, those who died from suicide in high-stigma areas were on average 18 years younger than those who died from suicide in low-stigma areas. This confirmed the findings of an earlier study that showed that lesbian, gay and bisexual youth in areas with high anti-gay prejudice were more likely to attempt suicide (Hatzenbuehler, 2011).

The authors of the earlier study pointed out similarities to other forms of minority status and structural stigma, and concluded that structural stigma also includes laws that criminalise, or restrict, the activities or identity of a minority group. One example are American laws that enforced racial segregation in some American states until the 1960s. A study that looked at the health consequences of structural stigma among Black people found that states with laws that enforced racial segregation had higher death rates of Black people (Krieger, 2012). Recent studies from the United States show that sexual orientation-related discriminatory laws and policies – laws and policies that deprive sexual minorities of certain rights (for example, the right to marry) – contribute to higher levels of mental health problems among sexual minority populations (Hatzenbuehler, Keyes and Hasin, 2009; Hatzenbuehler *et al.*, 2010). This is significant in the context of Southern and East Africa, where many countries have retained British colonial laws that criminalise consensual same-sex activity (Ambani, 2017), and thus discriminate against sexual and gender minority populations (Carroll and Mendos, 2017).


The findings that we present in this report demonstrate that, much like what we know from other contexts, sexual orientation and gender identity seem to be an influencing factor for people’s mental health and well-being, for their experiences of violence and for their access to healthcare.

3 For the purposes of this report, sexual minority people are those who do not identify as heterosexual, and are inclusive of the following: those who self-identify as lesbian, bisexual, gay, queer, pansexual, anyone who feels sexual attraction to, or has had sexual experiences with, a partner or partners of the same sex or gender, even if they self-identified as heterosexual, ‘men who have sex with men’ (MSM), and/or ‘women who have sex with women’ (WSW)

Similar to what researchers have observed in other parts of the world (Meyer, 2003; Mayer *et al.*, 2008b; Institute of Medicine, 2011b), we found disparities in health status between the LGBTI people participating in this study and data that exists for the general population: LGBTI people showed higher levels of mental health problems, experienced higher levels of violence and more barriers when accessing healthcare services. Drawing on the existing evidence on the impact of minority stress (Meyer, 2003) and structural stigma (Hatzenbuehler *et al.*, 2014), we argue that these disparities are due to the stigma, prejudice and social exclusion that LGBTI people experience due to their sexual orientation and/ or gender identity.

The structure of this report

This report consists of four sections. The first section is this introduction. The second section gives information about the methods we used in our study. We then move on to the third section to present our findings for the specific country under consideration: eSwatini. We first describe the socio-political context in which LGBTI people live. We then describe the research findings: first we describe the group of participants, then we describe the findings on health-seeking behaviour. We then describe the findings on experiences of violence, and after that describe the mental health outcomes of depression, anxiety, alcohol use, drug use, tobacco use and suicidality. When describing these findings, we compare our findings to what we know from studies with LGBTI people in other parts of the world, and to what we know about the general population in the specific country that the study was conducted in. Following this, we present an overview of the mental health outcomes for each specific population: for lesbian women, for gay men, for bisexual women and men, as well as for transgender people (including transgender women, transgender men and gender non-conforming people). This serves as an easy reference for anybody interested in population-specific health concerns. The fourth section of the report provides recommendations for governments, non-governmental organisations, academic researchers and international and national donors. In the appendices, we provide more detailed information about our methodology, and include the survey instrument.



METHODOLOGY

This section describes how we conducted the study. We explain how we planned the study, what questions we asked, and what we did with the data that we collected. We also provide details about who officially approved the study in the nine countries that we conducted it.

Participatory approach

For this study, we followed a community-based participatory research (CBPR) approach. Community-based research is a partnership approach to research that involves community members and academic researchers as partners in all stages of the research process. In this way, all partners can contribute their knowledge and skills, can decide jointly on what to research, how to do it, and what to do with the research findings. It also means that all partners share the responsibility and the ownership of the process and the research findings (Israel *et al.*, 1998).

CBPR is a well-used approach for studies that explore health-related disparities, particularly among marginalised communities, such as people of colour, or people living in poverty (Israel *et al.*, 2010). Because it directly involves communities as co-researchers, it is an excellent approach to examine the social context of health concerns (Leung, Yen and Minkler, 2004). Because it emphasises that power is shared between researchers and the community, and because it focuses on action based on the research findings, it also helps to minimise the understandable distrust of academic research that often exists among marginalised communities, who may see academics as mining information or misrepresenting them (Israel *et al.*, 2010).

The 23 community partner organisations for this study are listed in Table 1. The academic partner was the Gender Health and Justice Research Unit at the University of Cape Town in South Africa. Additional academic partners were Dr Chelsea Morroni from the Botswana UPenn Partnership and the Liverpool School of Tropical Medicine; Prof Adamson Muula from the College of Medicine, University of Malawi; Sindy Matse from the National AIDS Council in the Ministry of Health of eSwatini and Nelson Muparamoto from the University of Zimbabwe. The project was funded by COC Netherlands, who also provided logistical support throughout the process.

TABLE 1: Community partner organisations

Country	Partner Organisations
Botswana	
	Bonela
	LeGaBiBo
	Rainbow Identity Association
Ethiopia	
	<i>Names of the two organisations withheld for safety reasons</i>
Lesotho	
	The People's Matrix Association
Kenya	
	Ishtar-MSM
	Jinsiangu
	Maaygo
	Minority Womyn in Action
	National Gay and Lesbian Human Rights Commission (NGLHRC)
	Persons Marginalised and Aggrieved (PEMA)
Malawi	
	Centre for the Development of People (CEDEP)
South Africa	
	Durban Gay and Lesbian Community and Health Centre
	Gender Dynamix
	OUT LGBT Well-Being
	Triangle Project
Swaziland	
	The Rock of Hope
Zambia	
	Friends of Rainka
	Trans Bantu Zambia (TBZ)
	The Lotus Identity
Zimbabwe	
	Gays and Lesbians of Zimbabwe (GALZ)
	Sexual Rights Coalition (SRC)

Study design

Design of study aims

In October 2015, COC Netherlands held a consultative meeting with the community partner organisations and researchers from the Gender Health and Justice Research Unit (GHJRU) at the University of Cape Town. At that meeting, partner organisations identified the gaps in current research and knowledge on LGBTI people's health in the Southern and East African region. Additionally, the partner organisations, GHJRU researchers and COC discussed what study design would be best suited and discussed strategies for sampling and recruitment. These discussions identified a number of areas where more research was needed to better understand LGBTI health concerns. To address all of these areas was beyond the scope of this research project. We ranked all research needs that were identified and decided to focus on the top three: mental health and well-being, experiences of violence, and access to healthcare services.

Based on the discussions with the partner organisations, the GHJRU researchers drafted the study design. After all community partners, as well as COC Netherlands, provided feedback on our suggested study design, we finalised the study protocol and developed a survey questionnaire. Because there is currently little or even no research evidence on LGBTI people's mental health and well-being in our Southern and East African context, this project is an important opportunity to develop baseline data. For this reason, we developed a survey that could be used in all study countries, in order to compare findings across countries.

The survey

We reviewed national and international academic literature on how to measure mental health and well-being amongst LGBTI populations, specifically in Southern and East Africa. Based on these findings, we developed a draft for the survey we wanted to use in the study. We held two meetings with the community partner organisations and COC Netherlands to discuss the scope and wording of questions in the survey, and we revised the draft based on the feedback we received.

In each meeting, we held a group session to review the survey question by question and adjust the aims and wording of each section and question. As a team, we agreed to make small changes to standardised scales that measure mental health outcomes. While we wanted to create a single survey that could be used in all countries, in some instances we changed the wording of some of the questions for specific countries, so that participants would understand them better (for example, "apartment" versus "flat").

Once we had made all the suggested changes, we sent the survey to all community partner organisations and COC for a final round of feedback. Based on this last feedback, we finalised the survey.

Question design

All questions on the survey had categorical answers (answers that would organise participants into groups (categories), for example people who lived in Botswana, people who lived in Kenya, people who lived in South Africa, etc.). Only age, and number of cigarettes smoked per day were measured as continuous variables (information that can be measured on a scale or counted). For

many questions, we added an “Other, specify” option, so that participants could write or type additional/different information.

Socio-demographic measurement

We asked a number of questions to learn about participants’ socio-demographic circumstances. These included age, religion, education, housing, employment, race, and financial security (assessed by the question “On average do you have enough money to cover your basic needs?”). We created a variable to look at housing security, for which we asked participants if they owned their home, rented it, or shared a place with someone without paying rent. We classified participants who shared a place without financially contributing as ‘housing insecure’ because we hypothesised that they would be more vulnerable to being told to leave if their SOGIE was discovered by other people in the house. People who said they had no home, lived on the street, or lived in short-term accommodation (shelters) were also classified as housing insecure.

Measuring sexual orientation and gender identity

In public health literature, there is no recognised standard definition of sexual orientation or gender identity, nor is there consensus on how to measure them in quantitative studies. Sexual orientation is widely accepted as being comprised of three elements: sexual identity, sexual attraction, and sexual activity. A range of studies have used different combinations of these three elements to define participants’ sexual orientation (King *et al.*, 2008). In order to paint a nuanced picture of the participants’ sexual orientation, we aimed to assess each of these three elements.

1. **Sexual identity** was assessed by asking participants “In terms of your sexual orientation, how do you identify?” (Options: Lesbian, Bisexual, Gay, Heterosexual, Asexual, “Other, specify”)
2. **Attraction** was assessed by asking participants who they were sexually and emotionally attracted to (2 questions).
3. **Sexual activity** was assessed by asking participants about who they have had “sexual experiences with in the past year and their lifetime” (2 questions).

For attraction and sexual activity, the questionnaire gave participants a list of options from which they could select all that applied (Options: With women, with men, with trans women, with trans men, with gender non-conforming people, with intersex people, “I have not had sexual experiences”, “Other, specify”).

There is also no standardised way of asking participants about gender identity. We decided to combine three questions:

1. **Gender identity** was assessed by asking “In terms of your gender identity, how do you identify?” (Options: Woman, Man, Trans woman, Trans man, Gender non-conforming, “Other, specify”).
2. We asked about **sex assigned at birth** (Options: Male, Female, Intersex)
3. Additionally, we asked what sex/ gender was recorded in the participant’s identity document(s)

Based on participants' answers to these questions, we created categories for sexual orientation and gender identity. For sexual orientation, these were: lesbian, gay, bisexual, 'non-normative', and heterosexual. For gender identity, they were: cisgender women, cisgender men, transgender women, transgender men and gender non-conforming people. We use these categories to disaggregate the findings about experiences of violence and mental health outcomes. To create these categories, in some instances we had to re-code the way participants self-identified, based on the other information they provided in the questions about their sexuality and gender identity. The detailed algorithm for this re-coding is explained in Appendix 1.

Intersex participants

In our study, very few participants identified themselves as "intersex." Such small numbers make it difficult to draw statistical inferences about the data. For this reason, while the intersex participants are still included in the overall findings reported here, we do not disaggregate by intersex identity.

Measuring social support

We asked three questions about participants' social support: "Who do you go to when you need someone to talk to about problems in your life?", "Who in your life knows that you are LGBTI?", and "Of those, who have you told yourself about being LGBTI?" We combined the last two questions, to have an indicator of whether participants are 'out' in their social context.

Health-seeking behaviour and access to healthcare

We developed a number of general questions to ask about what kind of healthcare participants used, and where. Additionally, we adapted questions about experiences of discrimination in healthcare from other studies with LGBTI people (Bazargan and Galvan, 2012; Cruz, 2014; Calton, Cattaneo and Gebhard, 2015).

Measuring mental health and well-being

To measure depression and anxiety, as well as drug and alcohol use, we used internationally used and recommended scales. We chose scales that had been used in research on the African continent (specifically the countries in this study), and, if possible, that had been used in research with LGBTI people (anywhere in the world). However, there was little information about whether scales had been used with LGBTI populations (King *et al.*, 2008; Myer *et al.*, 2008; Chishinga *et al.*, 2011). We also considered the ease of understanding and potential ease of translation to other languages when choosing scales. Based on all these considerations, we used the following scales:

- The CES-D 10 (Center for the Epidemiological Studies of Depression Short Form) to measure depression. It is widely used to screen for signs of depression in primary care settings, and is often used for research on the prevalence of depression. It is important to keep in mind, however, that we cannot diagnose people using the CES-D 10. In order to receive a definitive diagnosis of clinical depression, an individual needs to see a healthcare provider.
- The Generalized Anxiety Disorder 7-item scale (GAD-7) to assess signs of anxiety that participants may have had in the last two weeks.
- The Alcohol Use Disorders Identification Test (AUDIT) to assess whether an participant's alcohol use is harmful.

- The Drug Use Disorders Identification Test (DUDIT) to assess if a participant's drug use is harmful.

To ask about suicide, we reviewed literature about LGBTI health to develop suicidality measures (Haas *et al.*, 2010; Marshall *et al.*, 2016).

In Appendix 1, we provide more detail on the scales and how we used the data we collected.

Measuring violence

We developed the questions that asked about experiences of violence based on the GHJRU's previous work in violence research. Additionally, we reviewed literature about intimate partner violence among LGBTI people (Calton, Cattaneo and Gebhard, 2015). We asked a series of "yes/no" questions about experiences with verbal harassment, emotional violence, physical violence ("Have you been physically assaulted?"), and sexual violence ("Have you been sexually assaulted?"). For physical and sexual violence, we asked about experiences in the last 12 months and in participants' lifetime. For participants who reported lifetime experiences of violence, we asked about three signs of post-traumatic stress based on the current *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5) of the American Psychiatric Association. These are: flashbacks or nightmares reliving the event; avoiding situation/people reminding them of the violent incident; jumpiness, irritability or restlessness following the incident (American Psychiatric Association, 2013).

Translations

The survey was translated into the following languages: Amharic, Chichewa, isiNdebele, Sesotho, Setswana, Shona, Siswati and Swahili. These translations were done by professional translators, and then reviewed by the community partner organisations. The changes that the partner organisations suggested were discussed with the professional translator, and incorporated into the final translated versions.

Fieldworker training

Each community partner organisation had a designated research coordinator and a research assistant. These two were responsible for training and overseeing fieldworkers, who collected data by handing out surveys to participants. We (the GHJRU researchers) trained the research coordinators and assistants in a three day 'Train the trainer workshop'. The training included information on research processes, how to make decisions about study design and methodology, best practices in data collection, research ethics and participant protection, as well as discussions about data analysis and the use of data once the study is over. We wrote a fieldworker manual, so that research coordinators and assistants would have the information from the training on hand. When organisations decided to employ additional fieldworkers, they were trained by the research coordinator.

Who could participate in the survey?

Eligibility to participate in the survey was defined by age, sexual orientation, and gender identity.

- **Be of adult age:** all participants needed to self-identify as being age 18 or older
- **Self-identified as LGBTI:** Participants were required to either not identify as heterosexual (and therefore be a sexual minority/member of the LGBTI community) or not be cisgender (and therefore be a gender minority, for example, transgender). Included in gender minorities are people with diverse sex characteristics (or who identified as intersex). We asked participants to self-identify. In the informed consent statement, we gave the following categorisations or identities as prompts to help potential participants determine their eligibility: gay, lesbian, bisexual, transgender, transsexual, transman, transwoman, intersex, queer, genderqueer, gender non-conforming, pansexual, omnisexual, men who have sex with men (MSM), women who have sex with women (WSW), kuchu.

Our study did not use a comparison group—that is, we did not survey people who identify strictly as heterosexual and cisgender. While this limits our ability to compare our findings about sexual and gender minority people with heterosexual and cisgender people, we draw on research with the general population to discuss possible differences between LGBTI people and heterosexual, cisgender people.

Sampling methodology

We combined two sampling methods to find research participants: community-based sampling and online-based sampling. This means that partner organisations would find participants at their events, or during their outreach activities, and also disseminate a link to an online version of the survey. In Appendix 1, we discuss in more detail why we chose these methods.

Neither of these two sampling methods allow us to draw inferences beyond the constituency population, meaning we will not be able to make predictions about larger LGBTI populations across the country or region. The findings from our study are therefore not representative of all LGBTI people in the participating countries, although they do give us an indication of what some of the problems affecting LGBTI people in these contexts maybe.

Each partner organisation aimed to enrol 200 participants. The numbers of participants in each country were therefore determined by the number of partner organisations in that country. In total, we analysed data from 3,796 participants. Table 2 shows the number of participants in each country. In Appendix 1, you will find a more detailed breakdown by country and organisation.

TABLE 2: Number of participants, by country

Country	Number of participants
Botswana	618
Ethiopia	198
Kenya	976
Lesotho	173
Malawi	197
South Africa	832
eSwatini	103
Zambia	353
Zimbabwe	346
TOTAL	3,796

Collecting data

As part of the participatory design of this project, each partner organisation designed an individual plan for recruiting participants, based on the recruitment plan that we have explained above. Organisations used a range of methods, including: promotion of the online survey through a facebook advert, promoting the survey among people who came for services at their office, recruiting through personal and professional networks of the fieldworkers.

The partner organisations used a mix of self-administration and fieldworker-administration to collect the data. **Self-administration** meant that the participant read the survey to themselves and filled it out on their own. **Fieldworker-administration** meant that a fieldworker read the questions to the participant.

Because questions about mental health, violence and experiences of discrimination might bring up traumatic memories or distress to people, all participants had access to psychosocial support, both during the data collection process and afterwards. In some organisations, this was provided by counsellors within the organisations, in others, through referrals to LGBTI-affirming counsellors outside of the organisation. All fieldwork teams held regular debriefing sessions for the fieldworkers, who also had access to the same psychosocial support services.

Pilot study

Before finalising the questionnaire, we conducted a pilot study in South Africa, the first country to implement data collection. The purpose of the pilot was to identify questions that should be added or removed, rephrased, or otherwise adjusted. The pilot study showed us a few questions that we needed to change in order to make the survey as easy to understand as possible. Once we made these changes, the questionnaire was considered final. We made no more changes to it during the study.

Analysing data

We entered all survey data into an online database called REDCap, an electronic data management system by Vanderbilt University, and then analysed it with the software Stata15. We ran descriptive statistics and measured associations between differences that we found among the participants in our sample. Where data was missing because participants had not answered a question, we used a method called ‘multiple imputation’.

For many key outcomes in this report, we report statistics for subgroups of the overall sample. We use this approach to highlight times when specific subgroups may be particularly vulnerable due to historical and persistent socio-economic disparities and oppression. However, we could only do this in countries where the size of the overall sample and subgroup were large enough to examine meaningfully.

Appendix 1 has more detailed information on our data analysis.

Research approvals and regulatory compliance

The study was approved by the University of Cape Town’s Faculty of Health Sciences Human Research Ethics Committee. Additionally, it was approved by national ethics or health regulatory bodies in each country (Table 3). In accordance with the guidelines for research on sexual and gender minorities’ health in rights-constrained environments and established best practices (amfAR, 2015; Amon *et al.*, 2012), in countries where obtaining regulatory approval would have significantly increased risks for our community partner organisations and/or research participants, we constituted a review board of community members to evaluate the risks and benefit of the study. This was overseen and approved by the University of Cape Town’s Faculty of Health Sciences Human Research Ethics Committee. We only enrolled participants who provided informed consent.

TABLE 3: Research approvals

Country	Approval authority	Reference number
Botswana	Review Board, Office of Research and Development, University of Botswana Ministry of Health and Wellness, Republic of Botswana	UBR/RES/IRB/ BIO/009 HPDME: 13/18/1
Ethiopia	Approval through community review board	-
Kenya	Kenya Medical Research Institute	KEMRI/RES/7/3/1
Lesotho	Research and Ethics Committee, Ministry of Health, Lesotho	ID94-2017
Malawi	University of Malawi, College of Medicine Research and Ethics Committee	P.01/18/2330
South Africa	University of Cape Town Faculty of Health Sciences Human Ethics Research Committee	HREC 012/2016
eSwatini	Scientific and Ethics Committee, Ministry of Health and Social Welfare, Kingdom of Swaziland	no reference number
Zambia	Approval through community review board	-
Zimbabwe	Medical Research Council of Zimbabwe	MRCZ/A/2303



FINDINGS IN ESWATINI

Sexual orientation, gender identity and expression in eSwatini

eSwatini, formerly the Kingdom of Swaziland, is an absolute monarchy landlocked between Mozambique and South Africa. It is one of the smallest countries on the African continent. eSwatini gained independence from Britain in 1968 and is ruled today by King Mswati III. Its history and economy is closely tied to neighbouring South Africa, but Swazis highly value its independent status, especially because of the years of apartheid in South Africa. The Swazi population is young: more than half of all Swazis are under the age of 20. Almost two thirds of Swazis live below the poverty line.

eSwatini's legal system, like that of many other former British colonies in Southern Africa, is derived from British colonial law and criminalises same-sex practices as 'sodomy', a common-law offence (Aengus Carroll and Mendos, 2017). In contradiction, Section 20 of the Swazi constitution guarantees all citizens the right to equality before the law (Kingdom of Swaziland, 2005).

The conflicting legal framework has led to stigma and marginalisation of sexual and gender minority Swazis. Whilst the Swazi government said that there have been no recent arrests under the colonial 'sodomy' law, the LGBTI advocacy organisations Rock of Hope and House Of Our Pride have documented numerous human rights violations ranging from violence to denial of public services (The Other Foundation, 2016). Sexual and gender minority Swazis are socially excluded, and a strong culture of silence around non-normative sexual orientations and gender identities means that they do not have access to psychosocial or other support structures. A recent report shows that many sexual and gender minority Swazis had been rejected by their families, and were without employment and stable housing (The Other Foundation, 2016).

What we know about the health and well-being of sexual and gender minority Swazis is largely limited to sexual health: we know that men who have sex with men have high risks for HIV and other sexually transmitted infections (Baral *et al.*, 2013). We know very little about the health and well-being of lesbian and bisexual women, or of transgender and gender non-conforming people. For men who have sex with men, we know little about their mental health and well-being (The Other Foundation, 2016).


The study population: sample characteristics

In eSwatini, we collected survey data on paper. Participants filled out surveys by themselves (self-administration) or with the assistance of a fieldworker (fieldworker-administration).

On paper, a total of 108 surveys were filled out through one partner organisation. Of these, 5 did not conform to the research protocol (protocol violations) because they either did not document informed consent or the respondent was not eligible (for example not 18 years old or older). We excluded these surveys from our final sample in analysis. We did not exclude any surveys because they were incomplete. A total of 103 paper surveys are included in the final sample.

Additionally, one response from the online survey is included in the eSwatini sample. During data collection in South Africa, a person living in eSwatini found the survey through facebook and participated. That response is included here, and not with the South African sample (Figure 1).

FIGURE 1: eSwatini participant sample



We do not report on the number of participants who were asked to participate, but who declined or were ineligible. These participants did not fill out the survey.

We therefore analysed 104 participant responses from eSwatini. Most of these (95%) filled out the survey on their own, and 5 participants (5%) filled it out with the help of a fieldworker.

Sociodemographic characteristics

Table 4 shows detailed information about participants' demographics (characteristics of the sample). The average (median) age was 26 years, with the youngest participant being 18 years old, and the oldest 52 years old. Less than half of participants lived in urban areas (43 participants, 41%), 39% lived in peri-urban areas (urban outskirts) (41), and 19% lived in a rural area (20). The majority of participants listed Christianity as their faith (84%).

TABLE 4: Sociodemographic characteristics

	n	%
Age group (n=86)		
18-24	29	33.72
25-34	46	53.49
35-44	9	10.47
45-54	2	2.33
Race (n=104)		
Black	101	97.12
White	1	0.96
Coloured	1	0.96
Other	1	0.96
What type of area do you live in? (n=104)		
Urban	43	41.35
Semi-urban/Peri-urban	41	39.42
Rural	20	19.23
Religious beliefs* (n=104)		
African tradition	3	2.88
Islam	2	1.92
Christianity	87	83.65
Not religious	12	11.54
Other, specify	1	0.96

*More than one answer possible

Sexual and gender diversity / sexual orientation and gender identity

Every participant in this study was a sexual minority and/or gender minority—because only people who identified as lesbian, gay, bisexual or any other non-heterosexual sexual orientation (sexual minorities), and/ or people who identified as transgender, gender queer, non-binary or any other non-cisgender gender identity (gender minorities) were allowed to participate in the survey. To determine participants' specific sexual orientations and gender identities, we asked a range of questions on sexual and emotional attraction, sexual behaviour, sexual identity, gender identity, sex classification at birth and legally assigned sex/gender. Participants' responses reflect the vast diversity of sexual and gender identity. We consider the diverse range of responses to these questions to be an important finding in itself, which highlights the diversity and complexity of sexual orientations and gender identities.

We first examined participants' self-identification of their sexual orientation and gender identity (see Table 5).

TABLE 5: Participant sexual orientation and gender identity, self-identified

Participant self-identified sexual orientation	Participant self-identified gender identity							
	Woman	Man	Trans woman	Trans man	GNC	Other	Missing data	Total
Lesbian	28	1	0	2	3	1	0	35
Bisexual	4	19	0	0	0	0	0	23
Gay	2	33	0	3	0	1	0	39
Heterosexual	0	0	1	0	0	0	0	1
'Other'	0	0	3	1	1	0	0	5
Missing data	0	0	0	1	0	0	0	1
Total	34	53	4	7	4	2	0	104

Table 5 describes how participants responded when asked how they identify their sexual orientation and gender identity, and therefore describes 'self-identification.' These are the terms that participants chose as most fitting to describe their sexual orientation and gender identity.

It should be noted that in Table 5, we did not categorise participants based on same-sex sexual experiences or the sex they were assigned at birth. Thus, Table 5 reflects only how people self-identified, and does not take into account, for example, people who identify as heterosexual but have had same sex/gender sexual relations, or who identify as man or woman, but were assigned a different sex at birth. In the 'other' category, four participants wrote that they identify their *sexual orientation* as 'transgender', which is widely understood to be a gender identity. One participant wrote that their sexual orientation was 'intersex'.

Throughout this report, we use categories of sexual orientation (lesbian, gay and bisexual) and gender identity (cisgender women, cisgender men, transgender women, transgender men and gender non-conforming people) to examine experiences of violence and mental health outcomes. To create these categories, we in some instances re-coded the way participants self-identified. This was to consider the additional information provided by other items in the survey. We describe the process of re-coding in the section 'Measuring sexual orientation and gender identity' in the previous section of this report.

Sexual minorities


We considered anyone who did not identify as heterosexual to be a sexual minority (see Table 5 and Figure 2), as well as anyone who felt sexual attraction to, or had had sexual experiences with, a partner or partners of the same sex or gender, even if they self-identified as heterosexual ($n=1$, 1%). In the existing HIV literature, these participants are referred to as 'men who have sex with men' (MSM), or 'women who have sex with women' (WSW) (Young and Meyer, 2005; Baral et al.,

2009). We decided to use the term sexual minority and not MSM or WSW for two reasons: (1) MSM and WSW are used in research on sexual behaviour and sexual health, and have been criticised for focusing too much on the sexual behaviour of people, while neglecting their relationships, communities and social networks; (2) the alternative term 'sexual minority' highlights people's social marginalisation due to non-normative sexual orientation or sexual practice. Given that our research is about people's mental health and well-being, and does not ask about sexual behaviour or sexual health, 'sexual minority' is more appropriate to highlight the effect of minority status on mental health, well-being, vulnerability to violence and marginalisation in healthcare.

In total, 97 participants (99%) were sexual minorities.

Figure 2 displays participants' sexual orientations. Participants who were classified as gay, lesbian and bisexual made up the majority of the sample. Additionally, one participant identified as heterosexual and five identified their sexual orientation as 'transgender' or 'intersex. When examining statistics by sexual orientation, we did not analyse the subgroup 'transgender' in comparison to the lesbian, gay and bisexual groups, because the number of people in it is too small to be statistically representative. However, these participants are still included in analysis of the overall sample.

Figure 2: Participants' sexual orientations



Gender minorities

In order to identify gender minority participants, we asked two questions: How did participants self-identify their gender identity (see Table 5), and what sex was assigned to participants at birth. Based on these parameters, we defined gender minority participants as:


- (1) those who self-identified as transgender women, transgender men, gender non-conforming (GNC) or other (15, 14%); and
- (2) those whose gender identity was different from the sex assigned to them at birth (n=5, 5%).

In total, 20 participants (19%) were gender minorities.

For this report, we considered those whose gender identity was different from the sex assigned to

them at birth to be transgender women and men, as appropriate. Figure 3 displays participants' gender identities.

FIGURE 3: Participants' gender identities



For more information about how we recorded and categorised sexual orientation and gender identity, see 'Sexual orientation and gender identity measurement' in the Methods of this report.

Socioeconomic circumstances

Table 6 details participants' socioeconomic status. For many key outcomes in this report, we report statistics for gender minority participants as a subgroup of the overall sample. We use this intersectional approach to highlight times when gender minority people, in comparison to cisgender people, may be particularly vulnerable due to stigma and persistent socio-economic disparities.

TABLE 6: Social and financial capital

	Overall sample (n=104)		Gender minority participants (n=20)		p
	n	%	n	%	
Housing type	(n=104)		(n=20)		
Categorical					<0.005*
House	52	50.00	17	85.00	
Apartment/flat	34	32.69	3	15.00	
Room	17	16.35	0	0.00	
Hut/traditional dwelling	1	0.96	0	0.00	
On the street	0	0.00	0	0.00	
Binary					0.023*
Informal	18	21.43	0	0.00	
Formal	66	78.57	20	100.00	

Overall sample (n=104)			Gender minority participants (n=20)		
	n	%	n	%	p
Housing security	(n=104)		(n=20)		0.011*
Owns home	12	11.54	1	5.00	
Rents home	50	48.08	5	25.00	
Shares housing without paying	42	40.38	14	70.00	

Highest completed level of education	(n=104)		(n=20)		0.645
No formal education	1	0.96	0	0.00	
Primary education	3	2.88	0	0.00	
Secondary school (matric)	43	41.35	7	35.00	
Post-secondary school/ University diploma or degree	57	54.81	13	65.00	

Employment	(n=102)		(n=20)		0.317
No employment	41	40.20	11	55.00	
Formal employment	39	38.24	6	30.00	
Informal employment	22	21.57	3	15.00	

Sufficient money for basic needs	(n=104)		(n=20)		0.640
No	62	59.62	11	55.00	
Yes	42	40.38	9	45.00	

Has medical aid	(n=91)		(n=19)		0.069
No	80	87.91	19	100.00	
Yes	11	12.09	0	0.00	

*Chi square/Fisher's exact test p-value significant, at $p < 0.05$

Seventy-nine percent of participants lived in housing or apartments (formal, stable housing structures). Of the others, most lived in 'rooms' (informal, unstable, or transient housing), and one lived in a hut/traditional dwelling. The sample was more diverse in respect of participants' housing security: almost half of participants rented (48%, 50 of 104 participants), another 40% lived in shared housing without paying rent (for example with family or friends; 40 of 104 participants) and 12% of participants owned their own home (12 of 104).

Levels of education were reported as high in the overall sample: 96% had completed secondary education (100 of 104 participants), and more than half of all participants had completed a post-secondary educational degree (for example, a tertiary degree or a post-secondary diploma; 57, 55%).

Many participants were in financially precarious situations: 40% did not have a paid job (41 out of 102 participants), and another 22% (22 of 102) held informal jobs, without contracts. Further, 60% (62 of 104) said they did not have enough money to cover their basic needs.

Only 12% of participants (11 out of 91) had private health insurance (medical aid).

Social support and being 'out'

To measure social support, we asked participants who they go to when they need to talk about life problems. We also asked who in their life knows about their sexual orientation and gender identity as a way of quantifying how 'out' they are. A description of these responses is in Table 7.

TABLE 7: Social support and being 'out'

	Overall sample (n=104)		Gender minority participants (n=20)		
	n	%	n	%	p
Who they go to for support	(n=103)		(n=20)		
Current partner(s)	59	57.28	9	45.00	0.216
Family member(s)	42	40.78	7	35.00	0.558
Friend(s)	77	74.76	15	75.00	0.978
Person/people living with	18	17.48	6	30.00	0.100
Healthcare provider(s)	16	15.53	3	15.00	0.941
Co-worker(s)	19	18.45	2	10.00	0.278
Person/people living nearby	2	1.94	1	5.00	0.270
LGBTI organisations	34	33.01	7	35.00	0.833

Who knows their SOGIE	(n=103)		(n=20)		
Current partner(s)	77	74.76	14	70.00	0.585
Family member(s)	61	59.22	12	60.00	0.937
Friend(s)	85	82.52	17	85.00	0.745
Person/people living with	32	31.07	12	60.00	0.002*
Healthcare provider(s)	31	30.10	6	30.00	0.992
Co-worker(s)	44	42.72	8	40.00	0.784
Person/people living nearby	28	27.18	7	35.00	0.381
LGBTI organisations	59	57.28	13	65.00	0.437

*Chi square/Fisher's exact test p-value significant, at $p < 0.05$

Participants had mostly disclosed their sexual orientation or gender identity to a friend (82%), to a current partner (75%) or to a family member (59%). Only one third of participants (30%) had disclosed their sexual orientation or gender identity to a healthcare provider or to people they lived with (31%). One in four participants (27%) had disclosed their sexual orientation or gender identity to people who lived nearby.

Overall, participants reported having the most social support from friends, current partners and family members and were 'out' to friends more than to other people in their lives.

Discrimination in healthcare

We asked participants about experiences of discrimination in health facilities, and how such experiences might have impacted their health-seeking behaviour. We examined experiences of discrimination or fear of discrimination in the overall sample and among gender minority participants. Table 8 describes these differences by these categories.

TABLE 8: Healthcare access and discrimination

	Overall sample (n=104)		Gender minority ⁴ participants (n=20)		
	n	%	n	%	p
Disclosed SOGIE to healthcare provider	(n=102)		(n=20)		
Yes	62	60.78	15	75.00	0.146
Has tried to hide SOGIE-related health concern from healthcare provider	(n=97)		(n=20)		
Yes	43	44.33	9	45.00	0.946
Have you been treated disrespectfully because of your SOGIE?	(n=104)		(n=20)		
Categorical	0.087				
Never	43	41.35	4	20.00	
Rarely	21	20.19	7	35.00	
Sometimes	31	29.81	6	30.00	
Often	9	8.65	3	15.00	
Binary	0.031*				
No (Never)	43	41.35	4	20.00	
Yes (Rarely/Sometimes/Often)	61	58.65	16	80.00	
Have you been called names or insulted in a health facility because of your SOGIE?	(n=104)		(n=20)		
Categorical	0.033*				
Never	61	58.65	10	50.00	
Rarely	24	23.08	2	10.00	
Sometimes	11	10.58	5	25.00	
Often	8	7.69	3	15.00	

4 Gender minority refers to all participants who were transgender, gender non-conforming or 'other' gender identities

	Overall sample (n=104)		Gender minority ¹ participants (n=20)		
	n	%	n	%	p
Binary					0.382
No (Never)	61	58.65	10	50.00	
Yes (Rarely/Sometimes/Often)	43	41.35	10	50.00	

Have you been denied healthcare because of your SOGIE?	(n=103)		(n=20)		
Categorical					0.068
Never	72	69.90	11	55.00	
Rarely	17	16.50	3	15.00	
Sometimes	11	10.68	4	20.00	
Often	3	2.91	2	10.00	
Binary					0.106
No (Never)	72	69.90	11	55.00	
Yes (Rarely/Sometimes/Often)	31	30.10	9	45.00	

*Chi square/Fisher's exact test p-value significant, at $p < 0.05$

More than half of participants (61%) had told a healthcare provider about their sexual orientation and/ or gender identity. Overall, the majority of participants had not been denied healthcare or called names or insulted due to their sexual orientation or gender identity. However, more than half of participants said that healthcare staff had treated them disrespectfully because of their sexual orientation or gender identity. Gender minority participants were significantly more likely to be treated disrespectfully. More than 40% of all participants had been insulted in a healthcare facility, and almost one third (30%) had been denied care because of their sexual orientation or gender identity. Participants' sexual orientation and gender identity also directly influenced healthcare, almost half (44%) of all participants reported trying to hide a health concern related to their sexual orientation or gender identity from a healthcare provider.

Recent research with lesbian, bisexual, gay and transgender Swazis confirms our findings that disrespectful and discriminatory treatment in health facilities is often directly linked to visible non-conforming sexual orientation or gender identity (Logie *et al.*, 2017, 2018). Previous research from eSwatini has also highlighted the discriminatory ways in which men who have sex with men are treated in HIV-related healthcare facilities, and how such experiences delay further health-seeking (Kennedy *et al.*, 2013). Our findings quantify these experiences and show that discrimination based on sexual orientation and gender identity and expression in Swazi healthcare facilities is widespread.

Health-seeking behaviour


We asked participants what health services they had sought in the previous year, and where they had gone for these services. Twelve percent of participants had private health insurance.

Overall, participants heavily relied on NGOs for health services. This is not surprising in light of the findings on stigmatisation and discrimination in healthcare facilities that we have described in the previous section.

Figure 4 shows health services that people had used in the previous year – NGOs, public healthcare facilities or private health care facilities. Overall, participants had most often gone to a health service when they were feeling sick, or for HIV voluntary testing and counselling (HIV VCT): 78% of participants (n=77) had gone for HIV VCT to community-based organisation in the past year, and 76% (n=74) had gone to public health facilities. 15% (n=15) of participants said that they had gone to a Community-based organization (CBO) for HIV care and treatment in the past year, and 21% (n=21) had gone to a public health facility. This suggests that the rate of participants living with HIV in our sample is at least between 15 and 21%.

Public health facilities were the most used source of care, except for sexual health preventive care (HIV VCT and barrier methods, for which participants mostly went to CBOs) and counselling and mental healthcare, for which participants also went to NGOs. Very few participants had accessed health facilities for gender affirming care. We explore this further in the following section.

FIGURE 4: Health-seeking behaviour in previous year



Gender affirming care

In addition to asking all participants about their general health-seeking behaviour, we also asked gender minority participants about their access to, and use of gender affirming practices. Participants' gender affirming practices are shown in Table 9. These findings are important because gender affirming practices such as binding⁵ are proven to support people's gender identity and expression, reduce psychological distress and increase their safety in public (Manderson 2012, Ekins and King 2006, Cole and Han 2011). However, some gender affirming practices also might have health implications (Peitzmeier et al. 2017). It is therefore important for NGOs and healthcare providers to know about the risks of gender affirming practices and to discuss them with people who want to use gender affirming practices, so that they can make informed choices and learn how to reduce these risks.

A third of gender minority participants who were assigned female at birth said that they used some form of binding (33%). Most of gender minority participants who were assigned male at birth said that they tucked (80%).

Fifteen percent of gender minority participants used hormones for gender affirmation. Because access to hormones is very limited (see the following paragraph), it is likely that some of those participants procure hormones through informal channels or outside of the country and use hormones without medical supervision.

TABLE 9: Gender affirming practices

Gender affirming practices among gender minority participants (n=20)		
	n	%
Binding (among those assigned female at birth, n=9)	3	33.33
Tucking (among those assigned male at birth, n=10)	8	80.00
Hormones (n=20)	3	15.00

While not all gender minority people need or desire gender affirming care, access also impacts the level of hormone use among gender minority participants. Therefore, Table 9 may not reflect the number of participants who want and need to use hormones but cannot access them. We asked participants who identified as transgender or gender non-conforming whether they had access to hormonal and surgical gender affirmation procedures through the local healthcare system (regardless of whether or not they wanted to actually make use of any of these). Table 10 shows that access to both hormonal and surgical gender affirmation was inaccessible: only one gender minority participant (8%) had access to hormone treatment, and no one had access to surgical procedures.

5 Binding is a technique to flatten one's breast or chest by using constrictive materials and clothing. Tucking is a technique to hide the bulge of male genitalia so that they are not conspicuous through clothing.

TABLE 10: Access to gender affirming care

Access to gender-affirming care for gender minority participants (n=20)		
	n	%
Access to hormones (n=13)	1	7.69
Access to surgical procedures (n=13)	0	0.00

Experiences of violence

We asked participants about their experiences of violence, including verbal harassment related to their sexual orientation and gender identity or expression (SOGIE) and experiences of physical violence, sexual violence and domestic violence. We asked about experiences of violence in the previous year, as well as at any point in participants' lifetime. Table 11 shows the findings for participants overall, and for gender minority participants.

Past research across the world has shown that LGBTI people are vulnerable to violence (Blondeel et al., 2018). In summary, our findings confirm these findings for eSwatini, where LGBTI people are vulnerable to violence, particularly physical violence. Vulnerability to violence is high across the different sexual orientations and gender identities.

TABLE 11: Experiences of harassment and violence

	Overall sample (n=104)		Gender minority participants (n=20)		p
	n	%	n	%	
SOGIE-related verbal harassment					
Experienced in lifetime	(n=102)		(n=20)		
	81	79.41	17	85.00	0.491
Experienced in past year	(n=89)		(n=18)		
	57	64.04	13	72.22	0.418
Sexual violence					
Experienced in lifetime	(n=103)		(n=20)		
	51	49.51	12	60.00	0.296
Experienced in past year	(n=103)		(n=20)		
	17	16.50	3	15.00	0.840

	Overall sample (n=104)		Gender minority participants (n=20)		p
	n	%	n	%	
Physical violence					
Experienced in lifetime	(n=103)		(n=20)		
	60	58.25	14	70.00	0.235
Experienced in past year	(n=102)		(n=20)		
	26	25.49	7	35.00	0.276

*Chi square/Fisher's exact test p-value significant, at $p < 0.05$


In our study, it is noteworthy that compared to cisgender participants, gender minority participants experienced higher levels of verbal harassment, physical and sexual violence.

In the following subsections, we discuss the different forms of violence (verbal, sexual and physical) in detail.

Verbal harassment

Seventy-nine percent of participants had experienced verbal harassment due to their sexual orientation and/or gender identity or expression at some point in their life, and 64% in the previous year (Figure 5). This number was even higher for gender minority participants: more than four out of five (85%) had experienced verbal harassment at some point in their life, and almost three in four (72%) in the previous year.


FIGURE 5: Verbal harassment, past year



Sexual violence

Half of all participants (50%) were survivors of sexual violence (Figure 6). This number was higher (60%) when we looked at only gender minority participants.

FIGURE 6: Sexual violence, lifetime




Of the overall participant group, almost one in six participants (17%) had experienced sexual violence in the previous year (Figure 7). Of gender minority participants, it was 15% of participants.


FIGURE 7: Sexual violence, past year



When disaggregated by sexual orientation and gender identity (see Figure 8 and Figure 9, and the tables on lesbian, bisexual and gay health in the Appendix), we found that half of the lesbian participants (51%) had experienced sexual violence in their lifetime (almost one in four, or 23% in the past year). Of all participants who identified as gay men, half (49%) had experienced sexual violence in their lifetime, and 21% in the previous year. Of all bisexual participants, 50% had experienced sexual violence in their lifetime, and none in the past year (note that the group of bisexual participants is quite small, therefore we could not disaggregate this data for bisexual women and bisexual men).

FIGURE 8: Sexual violence, by sexual orientation

Among gender minority participants, the levels of sexual violence were higher for lifetime sexual violence, compared with cisgender participants and the overall sample (see Figure 9): three in five (60%) gender minority participants had experienced sexual violence in their lifetime, compared to 48% of cisgender participants.

FIGURE 9: Sexual violence, by gender identity

The Bulletin of the World Health Organization recently published a systematic review on the prevalence of violence motivated by sexual orientation and/or gender identity (Blondeel et al., 2018). Its authors analysed data from 50 countries (the majority of data came from North America or Europe - data from African countries only came from Cote d'Ivoire and Rwanda). In

this systematic review, the prevalence of sexual violence experienced by lesbian, gay or bisexual people ranged from 2% to 10%. Our study findings show that the prevalence of sexual violence among lesbian, gay or bisexual people in eSwatini is significantly higher.

While we do not have data specifically from eSwatini, the South African Love Not Hate Campaign found that about one in ten (11%) of transgender participants had 'been sexually assaulted or raped' in the two previous years (OUT LGBT Well-being, 2016). This number is not disaggregated further, for example into trans women and trans men. Our findings suggest that gender minority people in eSwatini might face similar, or even higher levels of sexual violence. In the United States, the 2015 United States Transgender Survey showed that nearly half of transgender people (47%) have been sexually assaulted at some point in their lifetime, and one in ten (10%) have been sexually assaulted in the previous year (James *et al.*, 2016). Our findings suggest that the levels of sexual violence experienced by transgender people in eSwatini might be higher than in the US.


Our study did not collect data on the prevalence of sex work among participants. However, existing evidence shows that gender minority people are more likely to participate in sex work due to systemic, institutional and interpersonal discrimination that limits their access to education and work opportunities (Sausa, Keatley and Operario, 2007; Nadal, Davidoff and Fujii-Doe, 2014; The Other Foundation, 2016). For example, one in five participants (19%) in the 2015 United States Transgender Survey engaged in sex work for money, food, a place to sleep, or other goods or services (James *et al.*, 2016). In a South African study, transgender participants also spoke about exchanging sex with money or gifts during key informant interviews about access to sexual health services (Stevens, 2012). The higher risk of experiencing violence among sex workers, and the fact that gender minorities may be more likely to do sex work, may account in some part for the extremely high prevalence of sexual violence, as well as other forms of violence in our study.

It is quite likely, however, that our findings are still under-estimated – as recent research suggests (Müller and Meer, 2018), bias-motivated sexual violence against lesbian women can culminate in homicide. Because including data on deceased sexual and gender minority people was beyond the scope of our study, our findings might therefore underestimate the prevalence of bias-motivated sexual violence.


On the whole, our findings point out that a large number of LGBTI people in eSwatini, regardless of their specific sexual orientation or gender identity are survivors of sexual violence. The World Health Organization has shown that the health consequences of sexual violence are significant and diverse: they include physical injuries, unwanted pregnancy, sexually transmitted infections, including HIV, higher rates of mental health concerns, including depression and post-traumatic stress disorder, and higher likelihood of attempting suicide (Krug *et al.*, 2002). There is thus a need for LGBTI affirming counselling and psychosocial support, as well as medico-legal and court preparation services, should survivors decide to report and cases be brought to trial.

Physical violence

More than half of the participants in our study had experienced some form of physical violence at some point in their lives (Figure 10). More than two thirds of gender minority participants had experienced physical violence in their lifetime.

FIGURE 10: Physical violence, lifetime

In the year prior to answering the survey, one in four participants had experienced physical violence (25%, Figure 11). When examining gender minority participants, it was slightly more than one in three participants (35%).

FIGURE 11: Physical violence, past year

Compared by sexual orientation, gay and other men who have sex with men experienced the most physical violence: almost two thirds (64%) had experienced physical violence in their lifetime (Figure 12). Among lesbian and other women who have sex with women and bisexual participants, more than half said that they had experienced physical violence in their lifetime. In the past year, almost one in three lesbian women (29%), and one in four gay men (26%) had experienced physical violence.

FIGURE 12: Physical violence, by sexual orientation





Figure 13 shows that gender minority participants experienced even higher levels of violence, regardless of sexual orientation: more than two thirds of gender minority participants (70%) had experienced physical violence in their lifetime.

These violence levels are again much higher than what we know from existing studies, which are mostly from North America and Europe. Our findings show that 70% of gender minority participants had experienced physical violence in their lifetime. In the systematic review mentioned previously, studies show that between 3% and 31% of lesbian, bisexual and gay people said they had experienced any form of physical violence (Blondeel *et al.*, 2018). Among transgender people, it was 9% to 35%. In the United States, 27% of transgender people participating in a community-based survey said they had experienced physical violence in their lifetime (Bradford *et al.*, 2013). In a study among transgender women who have a history of sex work, also done in the United States, 51% of participants said they experienced physical violence in their lifetime (Nemoto, Bödeker and Iwamoto, 2011). Evidence suggests that transgender people are more vulnerable to violence if they experience more discrimination in their everyday lives (Bradford *et al.*, 2013). As our findings have shown, the gender minority participants in our study often live in financially precarious situations and have limited access to healthcare and socio-economic opportunities. While gender minority people worldwide may be more at risk for financial instability—for example, due to discrimination in employment—gender minority people in eSwatini may be particularly at risk due to the particular socioeconomic context. This might be one explanation why the levels of physical violence among our study’s transgender participants are much higher than in North America and Europe.

FIGURE 13: Physical violence, by gender identity

Perpetrators of violence

We asked participants who the perpetrators of violence were. Table 12 shows the details of this analysis. There are a few important observations, which we will describe in the following sections.

Intimate partner violence

First, we found high levels of intimate partner violence among participants. One in four participants (25%) said that they had been sexually assaulted by an intimate partner of any gender. Among lesbian participants, 29% had been sexually assaulted by an intimate partner. Among gay and other men who have sex with men, it was 26% - more than one in four (see also Table 15, Table 16, Table 17 and Table 20).

One in three participants had been physically assaulted by an intimate partner (34%). Among gay and other men who have sex men, it was almost half of participants (46%). Among gender minority participants, it was 40%.

Our study confirms findings from a representative national survey in the United States that found that levels of sexual and physical intimate partner violence are high among sexual minority men and women (Walters, Chen and Breiding, 2013).

TABLE 12: Perpetrators of lifetime sexual and physical violence

	Overall sample (n=104)		Gender minority participants (n=20)		p
	n	%	n	%	
Sexual violence					
Intimate partner	(n=102)		(n=20)		
	25	24.51	4	20.00	0.601
Someone known (not intimate partner)	(n=101)		(n=20)		
	28	27.72	6	30.00	0.799
Stranger	(n=100)		(n=19)		
	26	26.00	6	31.58	0.538
Someone lived with (intimate partner or other)	(n=98)		(n=20)		
	8	8.16	2	10.00	0.737
Physical violence					
Intimate partner	(n=103)		(n=20)		
	35	33.98	8	40.00	0.527
Someone known (not intimate partner)	(n=100)		(n=20)		
	34	34.00	6	30.00	0.673
Stranger	(n=98)		(n=20)		
	31	31.63	6	30.00	0.860
Someone lived with (intimate partner or other)	(n=99)		(n=19)		
	17	17.17	3	15.79	0.859
Participant felt any lifetime sexual or physical violence was linked to being LGBTI					
Yes	(n=68)		(n=16)		
	50	73.53	13	81.25	0.423

*Chi square/Fisher's exact test p-value significant, at $p < 0.05$

Stranger violence

Second, we found that strangers were often perpetrators of violence: one in three participants (32%) had experienced physical violence, and one in four participants (26%) sexual violence by strangers.

SOGIE-motivated violence

Third, almost three quarters of participants (74%), felt that the violence they experienced was linked to their sexual orientation and gender identity. Among gender minority participants, this was much higher: 81% of participants. While we cannot verify the motivation of the perpetrator(s), these findings contribute to the social context of violence motivated by sexual orientation or gender identity. Violence that is motivated by someone's sexual orientation or gender identity sends a message to all LGBTI people (Breen and Nel, 2011). This is detrimental to LGBTI people's mental health and well-being, as we will show in the coming sections of this report.

Impact of violence

We asked participants who had experienced sexual or physical violence in their lifetimes about three signs of post-traumatic stress. We classified participants who experienced all three symptoms as showing signs of post-traumatic stress. Half of the participants (51%) who had experienced violence showed signs of post-traumatic stress. This was the same among gender minority participants who had experienced violence (50%).

Reporting of violence

Participants who experienced any sexual or physical violence in the last year were asked about whether they reported it to the police, and if they had sought medical care (Table 13). About one in five of participants (21%) had reported to the police. More participants had gone to a healthcare provider for care (44%), though more than half had not.

TABLE 13: Reporting violence

Experienced violence in previous year	Overall sample (n=30)		Gender minority participants (n=7)		p
	n	%	n	%	
Sought medical care	(n=27)		(n=7)		
	12	44.44	2	28.57	0.326
Reported to police	(n=28)		(n=7)		
	6	21.43	1	14.29	0.595
Felt treated with less courtesy for being LGBTI	(n=10)		(n=1)		
Categorical					**
Never	3	30.00	0	0.00	
Rarely	0	0.00	0	0.00	
Sometimes	2	20.00	0	0.00	
Often	5	50.00	1	100.00	
Binary					**
No (Never)	3	30.00	0	0.00	
Yes (Rarely/Sometimes/Often)	7	70.00	1	100.00	

*Chi square/Fisher's exact test p-value significant, at $p < 0.05$; ** no test of significance performed due to small group numbers

Mental health outcomes

Mental health outcomes in the overall sample

Table 14 provides an overview of the mental health outcomes in the overall sample of participants. Additionally, the table also shows these mental health outcomes among all gender minority participants. Each of these health outcomes are described in further detail in the subsections below.

TABLE 14: Overall mental health outcomes

	Overall sample (n=104)		Gender minority ⁶ participants (n=20)		p
	n	%	n	%	
Depression (CES-D-10)	(n=103)		(n=20)		0.082
Classified as not depressed	54	52.43	7	35.00	
Classified as depressed	49	47.57	13	65.00	
Anxiety (GAD-7)	(n=101)		(n=19)		
Categorical					0.493
No signs of anxiety	49	48.51	7	36.84	
Signs of mild anxiety	36	35.64	8	42.11	
Signs of moderate anxiety	11	10.89	2	10.53	
Signs of severe anxiety	5	4.95	2	10.53	
Binary					0.495
No/mild anxiety	85	84.16	15	78.95	
Moderate/severe anxiety	16	15.84	4	21.05	
Suicidality					
Suicidal ideation, lifetime	(n=101)		(n=20)		
	47	46.53	11	55.00	0.397
Suicidal attempts, lifetime	(n=99)		(n=19)		
	26	26.26	8	42.11	0.081
Suicidal ideation, past year	(n=88)		(n=15)		
	12	13.64	3	20.00	0.430
Suicidal attempts, past year	(n=89)		(n=15)		
	6	6.74	2	13.33	0.264

⁶ Gender minority refers to all participants who identified as transgender, gender non-conforming or 'other' gender identities

	Overall sample (n=104)		Gender minority ⁶ participants (n=20)		p
	n	%	n	%	
Alcohol use	(n=97)		(n=19)		
Categorical					0.498
No alcohol use	33	34.02	9	47.37	
Some alcohol use	14	14.43	3	15.79	
Hazardous use	23	23.71	3	15.79	
Harmful use	13	13.40	3	15.79	
Alcohol dependence	14	14.43	1	5.26	
Binary					0.153
No/some alcohol use	47	48.45	12	63.16	
Hazard/Harm/ dependence	50	51.55	7	36.84	

Drug use	(n=98)		(n=19)		
Categorical					0.366
No drug use	78	79.59	14	73.68	
Some drug use	4	4.08	0	0.00	
Harmful drug use	15	15.31	5	26.32	
Drug dependence	1	1.02	0	0.00	
Binary					0.296
No/some drug use	82	83.67	14	73.68	
Harmful use/ dependence	16	16.33	5	26.32	

Tobacco use	(n=99)		(n=17)		0.778
Doesn't smoke at all	58	58.59	11	64.71	
Smoke some days	24	24.24	3	17.65	
Smoke everyday	17	17.17	3	17.65	

There is very limited data on the levels of mental health and well-being among the general population in eSwatini. For example, the most recent Demographic and Health Survey (Central Statistical Office and Macro International Inc., 2009) only asked about tobacco use, but no other mental health indicators. This makes it difficult to compare our findings to the levels of mental health and well-being of the general population. Regardless of what the levels of mental health concerns in the general Swazi population are, however, our findings show that among sexual and gender minority people, levels of depression, anxiety and substance use are high.


Depression

We used the instrument CES-D 10, a 10-item *Center for the Epidemiological Studies of Depression Short Form* to measure depression. It is widely used to screen for signs of depression in primary care settings, and is often used for research on the prevalence of depression. It is important to keep in mind, however, that we cannot diagnose people with the CES-D 10. In order to receive a definitive diagnosis of clinical depression, an individual needs to see a healthcare provider.


Based on the CES-D 10, almost half of our participants (48%) were classified as currently depressed (Figure 14).

During our literature search, we found little data on depression in eSwatini's general population for comparison. The latest Mental Health Atlas from eSwatini does not document prevalence of mental health treatment overall (World Health Organization, 2014a). The most recent available data was from 2003, in which 13% of men and 17% of women reported having received a depression diagnosis in the last 12 months (World Health Organization, 2003). A 2015 estimate from the Global Burden of Disease study put the prevalence of depressive disorders among all adults in eSwatini at 4.2% (World Health Organization, 2017a). This data does not disaggregate by sexual orientation and gender identity, and we need to assume that it was collected among a majority of heterosexual and cisgender people women and men. Compared to this data from the general population, our findings show that the levels of depression might be significantly higher among Swazi sexual and gender minority people.

FIGURE 14: Depression in overall sample



Although almost half of all participants were classified as depressed based on the CES-D 10 score, only one in nine participants (11%) said that they had previously been diagnosed with depression—fairly similar to the findings from the general population from 2003 (Figure 15) (World Health Organization, 2003). Of the 11 participants who had been diagnosed, three (27% of those diagnosed) were receiving treatment at the time of filling out the survey. When we looked at this in comparison to the participants' CES-D scores, 80% of those showing signs of depression had never been told by a healthcare provider that they have clinical depression. This suggests that there may be a large percentage of sexual and gender minority people who have not received diagnoses and treatment that could help them manage their symptoms of depression. As we saw in the findings on access to healthcare, very few participants in our study had accessed mental healthcare in the past year – this would explain why so few of participants with depression have not received treatment.

FIGURE 15: Depression diagnosis and treatment

When examining our participants' depression levels by sexual orientation and gender identity, we observed some differences. Lesbian and gay participants had much higher levels of depression than bisexual participants (see Figure 16). This was surprising, because in other study countries, bisexual participants did not have lower levels of depression. We think that this might be because there were very few bisexual women in our Swazi sample. In our study's South African sample, bisexual women had particularly high levels of depression. Because there were less bisexual women in our Swazi sample, it is possible that this lowered the overall percentage of bisexual people with depression.

A study with Swazi men who have sex with men (MSM), which asked whether they had "felt sad or had a depressed mood for more than 2 weeks at a time in the last 3 years", found that 65% said yes (Stahlman *et al.*, 2015). This question is a very different measure from the CES-D 10 that we used, but it confirms that feelings associated with depression may be prevalent. Qualitative research with MSM in eSwatini suggests that such feelings of depression are associated with feelings of shame from the social stigma of being sexually attracted to men, or from being diagnosed with HIV (Kennedy *et al.*, 2013).

Gender minority participants had higher level of depression (65%), compared to cisgender men (37%) and cisgender women (53%; Figure 17).

FIGURE 16: Depression, by sexual orientation





FIGURE 17: Depression, by gender identity




Based on the sparse available data from the general population data in eSwatini, it is difficult to assess whether levels of depression are higher among sexual and gender minority Swazis. Compared to the diagnosis rates in 2003, our reported level of diagnosis was similar. However, we also found that many more participants showed signs of depression based on the screening questions in CES-D 10. We could not find general population data using the CES-D 10 or a similar measure to assess depression. Given how limited access to mental healthcare is in eSwatini, the number of people who show symptoms of depression is probably much higher than the number of people who have been diagnosed with depression – simply because, in order to receive a diagnosis, one must see a healthcare provider. This is particularly true for sexual and gender minority people, who, as our findings show, have little access to healthcare facilities, and might avoid seeking care because they fear discriminatory treatment.

Anxiety


We used the instrument GAD-7 to assess signs of anxiety in participants, based on how they felt in the last two weeks. Based on the anxiety score (GAD-7), we classified participants into four categories: participants with no signs of anxiety, with signs of mild anxiety, with signs of moderate anxiety, and with signs of severe anxiety. The GAD-7 score should not be taken as a definitive diagnosis of anxiety in participants, but it is an assessment of current symptoms. According to the anxiety scores, just over half of participants (52%) had experienced anxiety in the last two weeks (see Figure 18). Five per cent of participants reported signs of severe anxiety.

FIGURE 18: Signs of anxiety, overall sample



We also asked participants if they had ever been diagnosed with anxiety. Overall, only 9% of participants said that they had previously been diagnosed by a healthcare worker with clinical anxiety. Of the 9 participants who had been diagnosed, only 4 (44%) were receiving treatment at the time of filling out the survey (Figure 19). Further, of the 15 participants who showed signs of anxiety, only 2 had been diagnosed with anxiety (13%). This suggests that sexual and gender minority people in eSwatini with anxiety symptoms (and possibly anxiety disorders) are not accessing the healthcare that they need.

FIGURE 19: Anxiety diagnosis and treatment




We did not find any actual comparison data for anxiety in eSwatini's general population. The latest Mental Health Atlas from eSwatini does not document prevalence of mental health treatment (World Health Organization, 2014a). We also did not find any literature on anxiety among Swazi sexual and gender minority people.

The Global Burden of Disease Study (2015) estimates the prevalence of anxiety disorders at 3.0% in eSwatini (World Health Organization, 2017b). Our findings suggest that anxiety may be more common among sexual and gender minority people in eSwatini.

We can draw on literature from other countries to help us understand our Swazi data. A recent systematic review found that most studies that looked at mental health indicate that sexual minority people show higher levels of anxiety than the general population (Plöderl and Tremblay, 2015). It is thus likely that the same is true for eSwatini – that our sexual and gender minority participants have higher levels of anxiety than the general population.


When we examined anxiety levels by sexual orientation, we saw differences by sexual orientation. Nearly a quarter of lesbian participants had moderate or severe signs of anxiety, meaning that an evaluation by a healthcare provider would be recommended. However, only 5% of bisexual participants had signs of moderate anxiety and no bisexual or gay participants showed severe anxiety (Figure 20, see also Table 15, Table 16 and Table 17).

FIGURE 20: Anxiety levels, by sexual orientation



When comparing anxiety levels by gender identity, we found that a similar amount of gender minority participants showed signs of anxiety, when compared to cisgender participants (see Figure 21, see also Table 20). However, while 10% of cisgender women and gender minority people showed signs of severe anxiety, no cisgender men showed severe signs.

FIGURE 21: Anxiety levels, by gender identity




Suicidality

We asked four questions about suicide: whether participants had thought about ending their life (suicidal ideation) at some point in their lives, and in the past year; and whether participants had tried to end their own life (suicide attempt) at some point in their lives, and in the past year (Table 14).


Figure 22 shows how many participants had ever thought about ending their life. Almost half of participants (47%) had thought about ending their life at least once at some point in their life. Of those who had thought about it, one in four (26%) had thought about ending their life in the previous year. Of all participants, 14% had thought of ending their life in the past year.

FIGURE 22: Suicidal ideation, overall sample




Approximately one in four participants (26%) had tried to end their life at some point in their lives. Seven percent had tried to end their life in the past year (Figure 23 and Table 14).

FIGURE 23: Suicide attempts, lifetime and past year




When looking at suicide attempts by participants of different sexual orientations (Figure 24, see also Table 15, Table 16 and Table 17) gay and other men who have sex with men had the highest level of attempted suicide (more than one in three attempted to end their life at some point; one in eight in the past year). Bisexual people had a 19% lifetime prevalence, and lesbian participants a 18% lifetime prevalence.

FIGURE 24: Suicide attempts, by sexual orientation



When comparing cisgender and gender minority participants (Figure 25, see also Table 20), we found that gender minority participants were more likely to attempt suicide: 42% of gender minority participants had attempted suicide at some point in their life, compared to 28% of cisgender men and 13% of cisgender women.


FIGURE 25: Suicide attempts, by gender identity

Examining the number of completed suicides among sexual and gender minority people in eSwatini was beyond the scope of this research and limits the interpretation of our findings. However, our findings highlight that it is vital to acknowledge that sexual and gender minority Swazis are particularly vulnerable to suicidality. Findings from other countries confirm our Swazi findings. A systematic review conducted by King and colleagues (2008) highlights the higher risk of suicidality that sexual minority people experience, though only studies from North America, Europe and Australasia were eligible to be included (further highlighting the need for research on the African continent). Their meta-analysis suggests that sexual minority people have about twice the risk of attempting suicide compared to non-sexual minorities (King *et al.*, 2008).

Alcohol use


We used the 10-item AUDIT instrument to ask participants about how much alcohol they consume, and about the impact of drinking alcohol on their lives. Figure 26 shows the levels of alcohol use in the overall sample. About a third of participants said they never drink alcohol (34%). Fifteen percent drank some alcohol without health risks. However, just over half of our participants (51%) drank alcohol at a level that had risks for their health: 37% showed signs of hazardous or harmful alcohol use, and 14% showed signs of alcohol dependence.

FIGURE 26: Alcohol use, overall sample




When looking at alcohol use by sexual orientation, we found that lesbian, bisexual, and gay participants had similar patterns of alcohol use. Bisexual participants showed the highest level of hazardous/harmful/dependent alcohol use (59%; for details, see Figure 27, Table 15, Table 16 and Table 17).

FIGURE 27: Alcohol use, by sexual orientation



When looking at alcohol use by gender identity (Figure 28 and Table 20), we found that cisgender participants had higher levels of hazardous/harmful/dependent drinking (54% for cisgender women and 56% of cisgender men) than gender minority participants (37%); however this difference was not statistically significant.

FIGURE 28: Alcohol use, by gender identity



Available data about alcohol consumption among the Swazi general population suggests that alcohol use, hazardous and otherwise, in our sample is high. The Swaziland STEPS Survey 2014, conducted among the general population, found that 73% of respondents had never drunk alcohol (Swaziland Government Ministry of Health, 2014). In our study with sexual and gender minority participants, it was only 34% who had never drunk alcohol. The Swaziland MICS Survey 2014, again conducted among the general population, describes the gender difference in alcohol consumption, with their findings suggesting that (cisgender) men are more likely to drink alcohol in their lifetime than (cisgender) women (76% of (cisgender) women reported never drinking as compared to 46% of men) (Central Statistical Office and UNICEF, 2016). However, older data from the WHO, published in 2010 is more in line with our findings, documenting that 30% of (cisgender) men and 57% of (cisgender) women had never drunk alcohol (again, compared to 34% overall in our study). WHO also found that 4% of “males” and 1% of “females” had alcohol dependence (World Health Organization, 2014b). Among our participants, 14% seemed to have alcohol dependence, which again suggests that alcohol use is higher among sexual and gender minority Swazis than among the general population.

The general population studies did not measure sexual orientation or detailed gender identity (beyond “men” and “women”), however, they did utilise strong sampling methodology which suggests they are representative findings at a population level.

At present there is a lack of data that is disaggregated by sexual orientation and gender identity in research on alcohol use (Flentje, Bacca and Cochran, 2015). International evidence on alcohol use among sexual minority people is somewhat mixed, although a 2008 systematic review shows that sexual minority people have higher levels of drinking than their heterosexual counterparts, and that sexual minority women may have more harmful use than sexual minority men (King *et al.*, 2008). It is unclear what motivates these differences or whether and how gender minority people were included in these studies. One qualitative study from eSwatini found that men who have sex with men (MSM) felt that MSM use alcohol as a coping mechanism against internalised stigma.

They believed alcohol is used to “gain confidence” prior to having sex—and, as a result, the sex they have may be less safe (Sithole, 2017).


In recent years, several new alcohol research studies have been conducted with gender minority people, though these have almost exclusively taken place in settings outside of the African continent. A few North American studies suggest that gender minority people are more likely to have harmful drinking practices than cisgender people, and that ‘gender minority stressors’ (Gonzalez et al 2017) may be associated with more alcohol use (Coulter et al., 2015; Scheim, Bauer and Shokoohi, 2016; Gonzalez, Gallego and Bockting, 2017).

Our findings confirm that in eSwatini, as in other settings, sexual and gender minority people have high levels of drinking. In comparison to general population data from eSwatini (not disaggregated by sexual orientation or gender identity) our sample of sexual and gender minority people shows higher levels of alcohol use.


Drug use

To measure levels of drug use among our sexual and gender minority sample, we used the DUDIT instrument (Figure 29). The majority of participants reported no drug use (80%), however, 15% reported harmful levels of drug use, including drug dependence (1%).

FIGURE 29: Drugs use, overall sample




When we examined differences between sexual orientation groups, lesbian participants reported the highest levels of drug use, and were the only group in which drug dependence was reported (1 participant). However, lesbian, gay and bisexual participants all reported similar levels of harmful/dependent drug use (18%, 14% and 13%, respectively; Figure 30).

FIGURE 30: Drug use, by sexual orientation

We observed some more notable differences in patterns of harmful drug use by gender identity (Figure 31). Gender minority participants in our sample had higher levels of harmful drug use (26%) than cisgender participants.

Findings from a study among cisgender men who have sex with men and transgender women in 8 African countries, including in eSwatini, supports our findings: 18% of their participants reported using some type of non-prescribed drug in the past 12 months (Poteat *et al.*, 2017). Drug use among our overall sample was higher (29%), but it is also difficult to compare findings in detail, due to different measures for drug use being used in different studies.

FIGURE 31: Drug use, by gender identity


As with alcohol, some research from other settings suggests that harmful drug use is more common among sexual and gender minority people than cisgender, heterosexual ones. Parental discomfort with homosexuality and being gender non-conforming in childhood have been found to be associated with higher substance-use rates (Rosario *et al.*, 2014).

Data from the WHO suggests that drug use in eSwatini is uncommon, estimating less than 1% of men and women have “drug use disorders” (World Health Organization, 2010). Thus, our findings show that sexual and gender minority people in eSwatini have higher levels of drug use than the general population.

Tobacco use


Forty-one percent of all participants reported that they smoke tobacco. Some smoke every day (17%) and some only on some days (24%) (Figure 32).

FIGURE 32: Tobacco use, overall sample




Overall, our findings suggest considerably higher levels of smoking among sexual and gender minority people, compared to the general population in eSwatini. The most recent Swaziland Demographic and Health Survey (DHS) (2006-07) documents that 2% of women and 22% of men in eSwatini use some form of tobacco. The DHS shows that as both men and women get older, they tend to use more tobacco (Central Statistical Office and Macro International Inc., 2009). More recently, the Swaziland STEPS Survey 2014 found that 1% of women and 12% of men smoke tobacco (Swaziland Government Ministry of Health, 2014). In our sample, participants of all sexual orientations and gender identities smoked more tobacco than those previous studies found in the general population.

In our sample, more than half of all lesbian participants (54%) smoked tobacco (see Figure 33, and also Table 15). One third of them smoked every day (33%). Among gay and other men who have sex with men, more than one third smoked (35%; see also Table 16). One in four smoked every day (24%).

FIGURE 33: Tobacco use, by sexual orientation

Among gender minority participants, more than one third smoked (36%; see Figure 34 and also Table 20). About one in five smoked every day.

FIGURE 34: Tobacco use, by gender identity

International data on smoking and sexual and gender minority people is limited. What is available, though mostly from Western countries, suggests that sexual and gender minority people have much higher rates of smoking tobacco than non-minorities (Blosnich, Lee and Horn, 2013; Lee *et al.*, 2014). Our findings confirm these international findings for sexual and gender minority people in eSwatini, who seem to smoke more than heterosexual, cisgender people in the general population.

Experiences of violence and health outcomes of lesbian participants

Lesbian participants include any person of any gender who self-identified their sexual orientation as 'lesbian', cisgender women who identified as 'gay' and transgender women who self-identified as 'gay' and had sex with or were attracted exclusively to women. There were 35 lesbian participants in the sample, including cis women, trans men and gender non-conforming people. No trans women in the eSwatini sample were categorised as lesbian (Figure 35).

FIGURE 35: Gender identities of lesbian participants




Table 15 shows the study findings for lesbian participants. Lesbian participants had some of the poorest health outcomes in our sample. Over half of lesbian participants were classified as depressed (63%), and one in four (24%) showed signs of moderate or severe anxiety. Lesbian participants had high use of drugs (27%) and alcohol (67%), much higher than available data on women in the general Swazi population (World Health Organization, 2010; Central Statistical Office and UNICEF, 2016). Four out of five (83%) said that they had been verbally harassed for their sexual orientation or gender identity, and over half had experienced physical violence and sexual violence (51% each). More than one in four had experienced sexual or physical violence by an intimate partner (29% and 29% respectively).

TABLE 15: Health outcomes and experiences of violence of lesbian participants

	n	%
Depression		
Depressed (based on CES-D 10) (n=35)	22	62.86
Ever been diagnosed with depression (n=33)	5	15.5
Of these, currently treated for depression (n=5)	1	20.00
Anxiety (n=33)		
Categorical		
Participants with no signs of anxiety	11	33.33
Participants with signs of mild anxiety	14	42.42
Participants with signs of moderate anxiety	4	12.12
Participants with signs of severe anxiety	4	12.12

	n	%
Binary		
No/mild anxiety	25	75.76
Moderate/severe anxiety	8	24.24
Ever been diagnosed with anxiety (n=32)	4	12.50
Of these, currently treated for anxiety (n=4)	0	0.00

Suicidality		
Suicidal ideation, lifetime (n=34)	13	38.24
Suicide attempt, lifetime (n=33)	6	18.18
Suicidal ideation, past year (n=30)	3	10.00
Suicide attempt, past year (n=30)	2	6.67

Alcohol use (n=33)		
Categorical		
No alcohol use	11	33.33
Some alcohol use	6	18.18
Hazardous use	7	21.21
Harmful use	5	15.15
Alcohol dependence	4	12.12
Binary		
No/some alcohol use	17	51.52
Hazard/Harm/ dependence	16	48.48

Drug use (n=33)		
Categorical		
No drug use	24	72.73
Some drug use	3	9.09
Harmful drug use	5	15.15
Drug dependence	1	3.03
Binary		
No/some drug use	27	81.82
Harmful use/ dependence	6	18.18

	n	%
Tobacco use (n=33)		
Don't smoke at all	15	45.45
Smoke some days	11	33.33
Smoke everyday	7	21.21
Verbal harassment for being LGBTI		
In lifetime (n=35)	29	82.86
Past year (n=29)	19	65.52
Sexual violence (n=35)		
In lifetime	18	51.43
Past year	8	22.86
Physical violence (n=35)		
In lifetime	18	51.43
Past year	10	28.57
Intimate partner, lifetime (n=35)		
Sexual violence	10	28.57
Physical violence	10	28.57

Experiences of violence and health outcomes of gay participants

Gay participants include all cisgender and transgender men who self-identified as gay; cisgender or transgender men who identified their sexual orientation 'MSM', as well as transgender women who self-identified as gay and were attracted to and had sex with men (transgender women who self-identified as gay but were exclusively attracted to or having sex with women were not included here—see the section on lesbian participants). There were 39 gay people in the sample. Figure 36 shows the gender identities of gay participants.

FIGURE 36: Gender identities of gay participants




Table 16 shows the study findings for gay participants. Over half of them were classified as depressed (51%), and one in eight showed signs of moderate or severe anxiety (13%). In this sample, gay participants had the highest levels of suicidality as compared to other sexual orientations: Two in five (39%) had attempted suicide in their lifetime, and one in eight (12%) in the past year. Half used alcohol in a harmful way (50%), and 13% used other drugs in a harmful way. Four out of five (84%) said that they had been verbally harassed for their sexual orientation or gender identity, over half (64%) had experienced physical violence, and almost half (49%) had experienced sexual violence in their lifetime. Almost half had experienced physical violence by an intimate partner, and one in four had experienced sexual violence by an intimate partner in their lifetime (46% and 26%, respectively).

TABLE 16: Health outcomes and experiences of violence of gay participants

	n	%
Depression		
Depressed (based on CES-D 10) (n=39)	20	51.28
Ever been diagnosed with depression (n=37)	5	13.51
Of these, currently treated for depression (n=5)	2	40.00

	n	%
Anxiety (n=39)		
Categorical		
Participants with no signs of anxiety	20	51.28
Participants with signs of mild anxiety	14	35.90
Participants with signs of moderate anxiety	5	12.82
Participants with signs of severe anxiety	0	0.00
Binary		
No/mild anxiety	34	87.18
Moderate/severe anxiety	5	12.82
Ever been diagnosed with anxiety (n=37)	4	10.81
Of these, currently treated for anxiety (n=4)	3	75.00

Suicidality		
Suicidal ideation, lifetime (n=39)	26	66.67
Suicide attempt, lifetime (n=38)	15	39.47
Suicidal ideation, past year (n=33)	8	24.24
Suicide attempt, past year (n=34)	4	11.76

Alcohol use (n=36)		
Categorical		
No alcohol use	14	38.89
Some alcohol use	4	11.11
Hazardous use	9	25.00
Harmful use	3	8.33
Alcohol dependence	6	16.67
Binary		
No/some alcohol use	18	50.00
Hazard/Harm/ dependence	18	50.00

Drug use (n=38)		
Categorical		
No drug use	32	84.21
Some drug use	1	2.63
Harmful drug use	5	13.16
Drug dependence	0	0.00

	n	%
Binary		
No/some drug use	33	86.84
Harmful use/ dependence	5	13.16

Tobacco use (n=38)		
Don't smoke at all	25	65.79
Smoke some days	9	23.68
Smoke everyday	4	10.53

Verbal harassment for being LGBTI (n=38)		
In lifetime	32	84.21
Past year	24	70.59

Sexual violence (n=39)		
In lifetime	19	48.72
Past year	8	20.51

Physical violence		
In lifetime (n=39)	25	64.10
Past year (n=38)	10	26.32

Intimate partner, lifetime		
Sexual violence (n=38)	10	26.32
Physical violence (n=39)	18	46.15

Experiences of violence and health outcomes of bisexual participants

Bisexual participants include any person who self-identified as bisexual. We provide overall numbers for all bisexual participants, as well as numbers for bisexual women (cis- and transgender women) and bisexual men (cis- and transgender men). Twenty-three participants in our sample were bisexual. As Figure 37 shows, most participants who identified as bisexual were cisgender men.

FIGURE 37: Gender identities of bisexual participants




Table 17 shows the study findings for all bisexual participants. Overall, bisexual participants tended to have somewhat better health outcomes than lesbian and gay participants. Because the number of cisgender and transgender women who identified as bisexual were too small to meaningfully include in statistical analysis, we have not disaggregated the findings by gender.

One in seven bisexual participants had signs of depression (14%), and 5% showed signs of moderate or severe anxiety. One in five (19%) had attempted suicide in their lifetime. More than half (59%) used alcohol at a level that might be harmful to their health, and one in seven (14%) used other drugs in a harmful way. One third (32%) used tobacco. Two in three (64%) said that they had been verbally harassed for their sexual orientation or gender identity, more than half (55%) had experienced physical violence, and half (50%) had experienced sexual violence. One in four had experienced sexual or physical violence by an intimate partner (24% and 24% respectively).

TABLE 17: Health outcomes and experiences of violence of bisexual participants

	All bisexual people (n=23)	
	n	%
Depression		
Depressed (based on CES-D 10) (n=22)	3	13.64
Ever been diagnosed with depression (n=21)	1	4.76
Of these, currently treated for depression (n=1)	0	0.00

	All bisexual people (n=23)	
	n	%
Anxiety (n=22)		
Categorical		
Participants with no signs of anxiety	15	68.18
Participants with signs of mild anxiety	6	27.27
Participants with signs of moderate anxiety	1	4.55
Participants with signs of severe anxiety	0	0.00
Binary		
No/mild anxiety	21	95.45
Moderate/severe anxiety	1	4.55
Ever been diagnosed with anxiety (n=21)	1	4.76
Of these, currently treated for anxiety (n=1)	1	100.00

Suicidality		
Suicidal ideation, lifetime (n=21)	5	23.81
Suicide attempt, lifetime (n=21)	4	19.05
Suicidal ideation, past year (n=20)	1	5.00
Suicide attempt, past year (n=19)	0	0.00

Alcohol use (n=22)		
Categorical		
No alcohol use	7	31.82
Some alcohol use	2	9.09
Hazardous use	6	27.27
Harmful use	4	18.18
Alcohol dependence	3	13.64
Binary		
No/some alcohol use	9	40.91
Hazard/Harm/ dependence	13	59.09


Drug use (n=21)		
Categorical		
No drug use	18	85.71
Some drug use	0	0.00
Harmful drug use	3	14.29
Drug dependence	0	0.00

	All bisexual people (n=23)	
	n	%
Binary		
No/some drug use	18	85.71
Harmful use/ dependence	3	14.29
Tobacco use (n=22)		
Don't smoke at all	15	68.18
Smoke some days	3	13.64
Smoke everyday	4	18.18
Verbal harassment for being LGBTI		
In lifetime (n=22)	14	63.64
Past year (n=20)	9	45.00
Sexual violence (n=22)		
In lifetime	11	50.00
Past year	0	0.00
Physical violence (n=22)		
In lifetime	15	54.55
Past year	3	13.64
Intimate partner violence (lifetime) (n=22)		
Sexual violence	5	22.73
Physical violence	5	22.73

Experiences of violence and health outcomes of gender minority participants

Gender minority participants include all participants who self-identified as transgender women, transgender men or gender non-conforming people. Additionally, it also includes all participants whose gender was different from the sex assigned at birth. Participants who selected 'other' gender identities and who were not cisgender are included in the overall number of gender minority people, but not reported as their own group due to their diversity and small numbers. There were 20 gender minority people in this sample. Figure 38 shows the sexual orientations of gender minority participants. The majority of gender minority people categorised as 'other' sexual orientation identified as heterosexual.

FIGURE 38: Sexual orientations of gender minority participants



Gender affirming care

It is worthwhile repeating the findings on gender affirming practices, as they relate directly to the health and well-being of transgender and gender non-conforming individuals. We asked gender minority participants about their access to and use of gender-affirming practices. To summarise the findings (detailed on page 35), a third of gender minority participants who were assigned female at birth said that they used binders (33%); and most of the gender minority participants who were assigned male at birth said that they tucked (80%). Fifteen percent of gender minority participants used hormones for gender affirmation.

TABLE 18: Gender affirming practices

Gender affirming practices among gender minority participants (n=20)		
	n	%
Binding (among those assigned female at birth, n=9)	3	33.33
Tucking (among those assigned male at birth, n=10)	8	80.00
Hormones (n=20)	3	15.00

Health outcomes

Table 19 shows the health outcomes for all gender minority people. Because the numbers in the specific sub-groups (e.g. transgender women, transgender men) were very small, we cannot report separately on these groups.

Two thirds of gender minority participants (65%) were classified as depressed, and one in five (21%) showed signs of moderate or severe anxiety. Over half (55%) had thought of suicide in their lifetime, and two in five (42%) had attempted suicide in their lifetime. One in eight (13%) had attempted suicide in the past year. Over one third (37%) used alcohol at levels that were harmful to their health; and one in four (26%) used drugs at harmful levels. One in three (35%) used tobacco. More than four in five (85%) had experienced verbal harassment due to their gender identity, almost two thirds (60%) had experienced sexual violence, and 70% had experienced physical violence. One in five had experienced either sexual violence (20%) by an intimate partner and two in five had experienced physical violence (40%) by an intimate partner.

TABLE 19: Health outcomes and experiences of violence of gender minority participants

	All gender minority people (n=20)	
	n	%
Depression		
Depressed (based on CES-D 10) (n=20)	13	65.00
Ever been diagnosed with depression (n=18)	1	5.56
Of these, currently treated for depression (n=1)	1	100.00
Anxiety (n=19)		
Categorical		
Participants with no signs of anxiety	7	36.84
Participants with signs of mild anxiety	8	42.11
Participants with signs of moderate anxiety	2	10.53
Participants with signs of severe anxiety	2	10.53
Binary		
No/mild anxiety	15	78.95
Moderate/severe anxiety	4	21.05
Ever been diagnosed with anxiety (n=18)	1	5.56
Of these, currently treated for anxiety (n=1)	0	0.00

	All gender minority people (n=20)	
	n	%
Suicidality		
Suicidal ideation, lifetime (n=20)	11	55.00
Suicide attempt, lifetime (n=19)	8	42.11
Suicidal ideation, past year (n=15)	3	20.00
Suicide attempt, past year (n=15)	2	13.33
Alcohol use (n=19)		
Categorical		
No alcohol use	9	47.37
Some alcohol use	3	15.79
Hazardous use	3	15.79
Harmful use	3	15.79
Alcohol dependence	1	5.26
Binary		
No/some alcohol use	12	63.16
Hazard/Harm/ dependence	7	36.84
Drug use (n=19)		
Categorical		
No drug use	14	73.68
Some drug use	0	0.00
Harmful drug use	5	26.32
Drug dependence	0	0.00
Binary		
No/some drug use	14	73.68
Harmful use/ dependence	5	26.32
Tobacco use (n=17)		
Don't smoke at all	11	64.71
Smoke some days	3	17.65
Smoke everyday	3	17.65
Verbal harassment for being LGBTI		
In lifetime (n=20)	17	85.00
Past year (n=18)	13	72.22

		All gender minority people (n=20)	
		n	%
Sexual violence			
In lifetime (n=20)		12	60.00
Past year (n=20)		3	15.00
Physical violence			
In lifetime (n=20)		14	70.00
Past year (n=20)		7	35.00
Intimate partner violence (n=20)			
Sexual violence		4	20.00
Physical violence		8	40.00

LIMITATIONS

This study has some limitations that should be kept in mind when reading the findings of this report.

First, because we recruited through organisations, we were likely to have participants who are already receiving some kind of services through these organisations. This means that the levels of mental health problems that we report might be higher than in a general sample of LGBTI people (Hendricks and Testa, 2012). We have tried to limit this potential over-estimation by also recruiting participants online, which in other studies has shown to reduce the over-estimation (Rosser *et al.*, 2007b). It is important to keep in mind, however, that even if the levels of mental health problems reported here are higher than among other LGBTI populations, they nevertheless present the current need for mental health support that our community partner organisations encounter through the services they offer.

Second, surveys that ask survivors of violence to report their experiences are likely to produce higher violence estimates than police-recorded administrative data. This is because often, violence is not reported to the police (which our findings confirm). Surveys with survivors of violence deal with incidents that not necessarily match the legal definition of a violent crime. Although data from surveys with survivors of violence are likely to elicit better disclosure of experiences of violence than data from police records, they can also be subject to undercounting, because some survivors may be reluctant to speak about their experiences. We have tried to reduce this potential under-estimation by collecting data through community partner organisations, with which many participants have a trustful relationship.

Third, we were faced with challenging decisions in how to categorise the diversity and complexity of sexual orientation and gender identity for the quantitative analysis. Based on the participatory methodology of this research, we used an in-depth discussion with South African partner organisations about the best way to do the categorisations. For example, a challenging decision was determining who should be included in the “lesbian” sexual orientation category. Although we considered categorising all transgender women who identified as gay to be “lesbian,” upon examination of these participants’ sexual behaviour and attraction, we noted that most gay transgender women strictly have sex with, and are attracted to, men. We therefore drew on sexual behaviour to make some coding decisions. We acknowledge that this may limit or bias our findings about sexual minority people. We have worked to describe our methodology openly to allow for interpretation and critique of these findings.

Fourth, this is an exploratory study. Neither of our two sampling methods allow us to draw inferences beyond the constituency population, meaning we are not able to make predictions about larger LGBTI populations across the country or region. The findings from our study are therefore not representative of all LGBTI people in the participating countries.

Last, it is difficult to compare findings on LGBTI people’s health across studies nationally and internationally. This is because there is currently no standardized measure of measuring or identifying sexual orientation and gender identity. As others have observed (Bradford *et al.*, 2013), the “lack of a standardized methodology to measure self-reported experiences of direct discrimination, lack of psychometric measures regarding validity or reliability of instruments,

potential reporting biases and measurement error, and variability in assessing chronic and acute exposures, as well as intensity, duration, and frequency of exposure” (Krieger, 1999) limit the current research evidence that we have on topics of discrimination and mental health.

CONCLUSION

Despite the limitations, our study is the first cross-sectional study to comprehensively describe levels of mental health among lesbian, gay, bisexual, transgender and intersex people in eSwatini. It shows that LGBTI people, regardless of their specific sexual orientation or gender identity, show higher levels of depression, anxiety, suicidality, and substance use than the general population. LGBTI people are also more likely to experience verbal harassment, physical and sexual violence than the general population, and face sexual orientation- and gender identity-related barriers when trying to access healthcare.

In 2014, the African Commission for Human and People’s Rights (ACHPR) passed Resolution 275, which calls for the protection from violence based on real or perceived sexual orientation and gender identity and proposes specific obligations for African states (ACHPR, 2014). At a joint dialogue of the ACHPR, the Inter-American Commission on Human Rights and the UN, participants concluded that: “[d]ata and evidence is critical to understand the extent and gravity of violations and to advocate for the adoption of measures to prevent, address and redress human rights violations faced by [sexual and gender minorities]” (ACHPR, 2016). The findings from our study provide such data for eSwatini, and evidence the seriousness of the rights violations against Swazis who identify as sexual or gender minorities, as well as the health consequences.

The findings from our study confirm that in eSwatini, as described in other parts of the world (Meyer, 2003; Hatzenbuehler *et al.*, 2014), social exclusion, marginalisation and stigma due to non-normative sexual orientation and/ or gender identity has a negative impact on the mental health and wellbeing of people who identify as lesbian, gay, bisexual, transgender or intersex.

The findings from our study demonstrate the urgent need for mental health services that are affirming of sexual and gender diversity and are provided without sexual orientation and gender identity-related stigma, prejudice and discrimination. It is clear that affirming and non-judgmental mental healthcare services for sexual and gender minority people are at least as important as HIV-related health services. This is not just to improve mental health and wellbeing, but also to support efforts to decrease the vulnerability to HIV. Our findings provide important contextual information to healthcare providers on the health impact of social exclusion based on sexual orientation and/ or gender identity.

In summary, our report paints a sobering picture of the state of mental health and well-being of LGBTI people in eSwatini. It underscores the responsibilities that government, NGOs, researchers and funders have to address both the health concerns and its underlying causes. These underlying causes are rooted in the criminalisation of same-sex activity, in stigma, discrimination and marginalisation, and in prejudicial and biased attitudes by healthcare providers, other civil servants and the general population.

RECOMMENDATIONS

Recommendations for national government

- Decriminalise same-sex activity: legal reform to abolish laws which contribute to sexual orientation and gender identity-related stigma, prejudice and discrimination against sexual and gender minority people living in eSwatini, including men who have sex with men and women who have sex with women.
- Take into account sexual and gender diversity when programming for gender issues, including gender-based violence;
- Improve access to mental health services for LGBTI populations:
 - Ensure that mental health services are affirming of sexual and gender diversity;
 - Ensure that mental health services are provided without sexual orientation and gender identity-related stigma, prejudice and discrimination;
 - We recommend following the guidelines on sexual and gender diversity published by the Psychological Association of South Africa;
 - Include mental health assessments, care and referrals into the HIV-related package of care for key populations.
- Build knowledge, skills and capacity within the public health sector to reduce sexual orientation and gender identity-related stigma, prejudice and discrimination in healthcare:
 - Provide mandatory sensitisation on sexual orientation, gender identity and expression, as well as values clarification, for healthcare providers at health facilities;
 - Provide continuous professional development education and training for healthcare providers to raise awareness of the mental health needs of LGBTI people in eSwatini;
 - Include teaching on sexual orientation and gender identity-related health concerns into health professions education.
- Support the work of civil society organisations who provide services, including mental healthcare, for sexual and gender minorities.

Recommendations for civil society organisations

- For LGBTI civil society organisations:
 - Provide affirming counselling services for LGBTI people, and actively raise funds for such services;
 - Recognise that staff at LGBTI civil society organisations may have experiences with violence, or mental health concerns, and prioritise interventions and programmes for staff well-being;
 - Include mental health as an important aspect of the health of LGBTI people in advocacy, programming and outreach work;
 - Build relationships and referral services with mental healthcare providers who are

willing to provide LGBTI-affirming services.

- Continue advocacy, public awareness and values clarification work to address the causes of violence, namely discrimination, stigma and prejudicial social and cultural attitudes.
- For civil society organisations providing services to survivors of violence:
 - Ensure that all staff, especially psychosocial and court support staff, are able to provide affirming services to LGBTI survivors of violence;
 - In gender-based violence advocacy and programming, take into account how sexual orientation, gender identity and expression can increase vulnerability to gender-based violence;
 - Actively build links to LGBTI civil society organisations.

Recommendations for donors

- Provide funding for services, programming and advocacy work linked to mental health and sexual orientation, gender identity and expression;
- Raise awareness of the need for mental health services and education for LGBTI people with other donors;
- Ensure that funds for violence prevention and programming build programmes that take into account vulnerabilities linked to sexual orientation, gender identity and expression, and are inclusive of people with diverse sexual orientations and gender identities and expressions.

Recommendations for academics and researchers

- Work with civil society organisations to establish research priorities and thematic areas, and fully and meaningfully involve civil society organisations in research projects:
- Follow existing guidelines on how to work with LGBTI populations in health-related research, for example the *Guidelines for Conducting Participatory Social Research with Key Populations and Marginalised Communities* (KP Reach, 2018).
 - Meaningfully include civil society organisations in the development of research proposals, including in budget items.
- Include demographic data on sexual orientation and gender identity and expression in population-based studies, in order to expand the knowledge base on sexual orientation, gender identity and expression and health.
- Conduct research, in partnership with civil society organisations, to further understand the mental health and well-being of LGBTI populations in eSwatini.

REFERENCES

- Academy of Science of South Africa (2015) *Diversity in Human Sexuality: Implications for Policy in Africa*. Pretoria. Available at: <https://www.assaf.org.za/files/8-June-Diversity-in-human-sexuality1.pdf>.
- African Commission on Human and Peoples' Rights (2014). *275: Resolution on Protection against Violence and other Human Rights Violations against Persons on the basis of their real or imputed Sexual Orientation or Gender Identity*. Banjul, Gambia: ACHPR.
- African Commission on Human and Peoples' Rights (2016). *Ending violence and other human rights violations based on sexual orientation and gender identity: A joint dialogue of the African Commission on Human and Peoples' Rights, Inter-American Commission on Human Rights and United Nations*. Pretoria: ACHPR.
- Ambani, J. O. (2017) 'A triple heritage of sexuality? Regulation of sexual orientation in Africa in historical perspective', in Namwase, S. and Jjuuko, A. (eds) *Protecting the Human Rights of Sexual Minorities in Contemporary Africa*. Pretoria: Pretoria University Law Press (PULP), pp. 14–51.
- American Psychiatric Association (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: American Psychiatric Publishing.
- amfAR, International AIDS Vaccine Initiative, Johns Hopkins University - Center for Public Health and Human Rights, United Nations Development Programme (2015). *Respect Protect Fulfill: Best practices guidance in conducting HIV research with gay, bisexual, and other men who have sex with men (MSM) in rights- constrained environments*. amfAR.
- Amon, J.J., Baral, S.D., Beyrer, C., Kass, N.E. (2012). 'Human Rights Research and Ethics Review: Protecting Individuals or Protecting the State?' *PLOS One*, 9:e1001325.
- Baral, S. et al. (2009) 'HIV prevalence, risks for HIV infection, and human rights among men who have sex with men (MSM) in Malawi, Namibia, and Botswana.', *PloS one*, 4(3), p. e4997.
- Baral, S. D. et al. (2013) 'A cross-sectional assessment of the burden of HIV and associated individual- and structural-level characteristics among men who have sex with men in Swaziland.', *Journal of the International AIDS Society*, 16 Suppl 3, p. 18768.
- Barbor, T. F. et al. (2001) 'AUDIT: The Alcohol Use Disorders Identification Test: Guidelines for Use in Primary Care. Second Edition.' Geneva: World Health Organization Department of Mental Health and Substance Dependence.
- Bazargan, M. and Galvan, F. (2012) 'Perceived discrimination and depression among low-income Latina male-to-female transgender women.', *BMC public health*. BioMed Central, 12, p. 663.
- Berman, A. H. et al. (2003) 'DUDIT The Drug Use Disorders Identification Test Manual'. Stockholm: Karolinska Institutet, Department of Clinical Neuroscience, Section for Alcohol and Drug Dependence Research.
- Bhugra, D. et al. (2016) 'WPA Position Statement on Gender Identity and Same-Sex Orientation, Attraction and Behaviours', *World Psychiatry*, 15(3), pp. 299–300.
- Blondeel, K. et al. (2018) 'Violence motivated by perception of sexual orientation and gender identity: a systematic review', *Bulletin of the World Health Organization*, 96(1), p. 29–41L.
- Blosnich, J., Lee, J. G. L. and Horn, K. (2013) 'A systematic review of the aetiology of tobacco disparities for sexual minorities.', *Tobacco control*. NIH Public Access, 22(2), pp. 66–73.
- Bockting, W. O. et al. (2013) 'Stigma, Mental Health, and Resilience in an Online Sample of the US Transgender Population', *American Journal of Public Health*, 103(5), pp. 943–951.
- Bradford, J. et al. (2013) 'Experiences of transgender-related discrimination and implications for health: Results from the Virginia transgender health initiative study', *American Journal of Public Health*, 103(10), pp. 1820–1829.
- Breen, D. & Nel, J.A., 2011. South Africa - A home for all? The need for hate crime legislation. *South African Crime Quarterly*, (38), pp.33-43.
- Calton, J.M., Cattaneo, L.B. & Gebhard, K.T., 2015. Barriers to Help Seeking for Lesbian, Gay, Bisexual, Transgender, and Queer Survivors of Intimate Partner Violence. *Trauma, Violence & Abuse*, 17(5), pp. 585-600.
- Carroll, A. and Mendos, L. R. (2017) *State-Sponsored Homophobia - A world survey of sexual orientation laws: Criminalisation, protection and recognition*. Geneva: ILGA. Available at: http://ilga.org/downloads/02_ILGA_State_Sponsored_Homophobia_2016_ENG_WEB_150516.pdf.
- Central Statistical Office and Macro International Inc. (2009) *Swaziland Demographic and Health Survey 2006-07*. Mbabane. doi: 10.1111/j.1728-4465.2009.00189.x.
- Central Statistical Office and UNICEF (2016) 'Swaziland Multiple Indicator Cluster Survey 2014. Final Report'. Mbabane, Swaziland: Central Statistical Office and UNICEF. Available at: http://www.swazistats.org.sz/images/reports/Multiple_Indicator_Cluster_Survey_2014.pdf.

- Chishinga, N. et al. (2011) 'Validation of brief screening tools for depressive and alcohol use disorders among TB and HIV patients in primary care in Zambia', *BMC Psychiatry*, 11(1), p. 75.
- Cole, B., and L. Han. (2011) *Freeing Ourselves: A Guide to Health and Self Love for Brown Bois*. Oakland: Brown Boi Project.
- Coulter, R. W. S. et al. (2015) 'Differences in alcohol use and alcohol-related problems between transgender- and nontransgender-identified young adults', *Drug and Alcohol Dependence*, 154, pp. 251–259.
- Cruz, T. M. (2014) 'Assessing access to care for transgender and gender nonconforming people: A consideration of diversity in combating discrimination', *Social Science & Medicine*, 110, pp. 65–73.
- De Cuypere, G. and Winter, S. (2016) 'A gender incongruence diagnosis: where to go?', *The Lancet Psychiatry*, 3(9), pp. 796–797.
- Drescher, J. (2015) 'Queer diagnoses revisited: The past and future of homosexuality and gender diagnoses in DSM and ICD', *International Review of Psychiatry*, 27(5), pp. 386–395.
- Ekins, R., and D. King. (2006) *The Transgender Phenomenon*. Thousand Oaks, CA: SAGE.
- Flentje, A., Bacca, C. L. and Cochran, B. N. (2015) 'Missing data in substance abuse research? Researchers' reporting practices of sexual orientation and gender identity', *Drug and Alcohol Dependence*, 147, pp. 280–284.
- Frost, D. M. and Meyer, I. H. (2009) 'Internalized Homophobia and Relationship Quality among Lesbians, Gay Men, and Bisexuals.', *Journal of counseling psychology*, 56(1), pp. 97–109.
- GATE (2017) *Submission by GATE to the World Health Organization : Intersex codes in the International Classification of Diseases (ICD) 11 Beta Draft*. New York: GATE.
- Gergen, K. J. (2001) 'Psychological science in a postmodern context', *American Psychologist*, 56(10), pp. 803–813.
- Gonzalez, C. A., Gallego, J. D. and Bockting, W. O. (2017) 'Demographic Characteristics, Components of Sexuality and Gender, and Minority Stress and Their Associations to Excessive Alcohol, Cannabis, and Illicit (Noncannabis) Drug Use Among a Large Sample of Transgender People in the United States', *The Journal of Primary Prevention*, 38(4), pp. 419–445.
- Haas, A. P. et al. (2010) 'Suicide and Suicide Risk in Lesbian, Gay, Bisexual, and Transgender Populations: Review and Recommendations', *Journal of Homosexuality*, 58(1), pp. 10–51.
- Hatzenbuehler, M. L. et al. (2010) 'The impact of institutional discrimination on psychiatric disorders in lesbian, gay, and bisexual populations: A prospective study', *American Journal of Public Health*, 100(3), pp. 452–459.
- Hatzenbuehler, M. L. (2011) 'The Social Environment and Suicide Attempts in Lesbian, Gay, and Bisexual Youth', *Pediatrics*, 127(5), pp. 896–903.
- Hatzenbuehler, M. L. et al. (2014) 'Structural stigma and all-cause mortality in sexual minority populations', *Social Science and Medicine*, 103, pp. 33–41.
- Hatzenbuehler, M.L. et al., 2018. Corrigendum to 'Structural stigma and all-cause mortality in sexual minority populations'; [Soc. Sci. Med. 103 (2014) 33-41]. *Social Science & Medicine*, 200, p.271.
- Hatzenbuehler, M. L., Keyes, K. M. and Hasin, D. S. (2009) 'State-level policies and psychiatric morbidity in lesbian, gay, and bisexual populations', *American Journal of Public Health*, 99(12), pp. 2275–2281.
- Hendricks, M. L. and Testa, R. J. (2012) 'A conceptual framework for clinical work with transgender and gender nonconforming clients: An adaptation of the minority stress model', *Professional Psychology: Research and Practice*, 43(5), pp. 460–467.
- Herek, G. M., Gillis, J. R. and Cogan, J. C. (2009) 'Internalized stigma among sexual minority adults: Insights from a social psychological perspective.', *Journal of Counseling Psychology*, 56(1), pp. 32–43.
- Institute of Medicine, 2011. *The Health of Lesbian, Gay, Bisexual, and Transgender People: Building a Foundation for Better Understanding*. Washington DC: Institute of Medicine.
- Israel, B. A. et al. (1998) 'Review of Community-Based Research: Assessing Partnership Approaches to Improve Public Health', *Annual Review of Public Health*, 19(1), pp. 173–202.
- Israel, B. A. et al. (2010) 'Community-based participatory research: A capacity-building approach for policy advocacy aimed at eliminating health disparities', *American Journal of Public Health*, 100(11), pp. 2094–2102.
- James, S.E. et al., 2016. *The Report of the 2015 U.S. Transgender Survey*, Washington, DC: National Center for Transgender Equality.
- Kara, S., 2017. *Gender is not an illness: How pathologizing trans people violates international human rights law*. Berlin: GATE.
- Kennedy, C. E. et al. (2013) "'They are human beings, they are Swazi": intersecting stigmas and the positive health, dignity and prevention needs of HIV-positive men who have sex with men in Swaziland.', *Journal of the International AIDS Society*, 16 Suppl 3, p. 18749.
- King, M. et al. (2008) 'A systematic review of mental disorder, suicide, and deliberate self harm in lesbian, gay and bisexual people', *BMC*

- Psychiatry*. BioMed Central, 8(1), p. 70. doi: 10.1186/1471-244X-8-70.
- Kingdom of Swaziland (2005) *The Constitution of the Kingdom of Swaziland Act 2005*.
- KP Reach (2018). 'Guidelines for Conducting Participatory Social Research with Key Populations and Marginalised Communities'. Available online at: <https://kpreach.net/resources/#guidelines> (accessed 11 May 2019).
- Krieger, N., 2012. Methods for the scientific study of discrimination and health: An ecosocial approach. *American Journal of Public Health*, 102(5), pp.936–945.
- Krieger, N., 1999. Embodying inequality: a review of concepts, measures, and methods for studying health consequences of discrimination. *International Journal of Health Services*, 29(2), pp.295-352.
- Kroenke, K., Spitzer, R. L. and Williams, J. B. (2001) 'The PHQ-9: validity of a brief depression severity measure.', *Journal of general internal medicine*. Springer, 16(9), pp. 606–13. doi: 10.1046/J.1525-1497.2001.016009606.X.
- Krug, E. E. G. et al. (2002) *The world report on violence and health*, World Health Organization. Geneva. doi: 10.1136/ip.9.1.93.
- Lee, J.G.L. et al., 2014. Promotion of tobacco use cessation for lesbian, gay, bisexual, and transgender people: a systematic review. *American Journal of Preventive Medicine*, 47(6), pp.823–31.
- Leung, M. W., Yen, I. H. and Minkler, M. (2004) 'Community-based participatory research: A promising approach for increasing epidemiology's relevance in the 21st century', *International Journal of Epidemiology*, 33(3), pp. 499–506.
- Logie, C. (2012) 'The case for the World Health Organization's Commission on the Social Determinants of Health to address sexual orientation.', *American Journal of Public Health*, 102(7), pp. 1243–6.
- Logie, C. et al. (2017) 'Barriers and facilitators to engagement in the HIV prevention cascade among lesbian, gay, bisexual, and transgender persons in Swaziland', *Sexually Transmitted Infections*, 93(Suppl 2), p. A19.
- Logie, C. et al. (2018) 'Marginalization and social change processes among lesbian, gay, bisexual and transgender persons in Swaziland: implications for HIV prevention', *AIDS Care*. Taylor & Francis, 30(sup2), pp. 33–40.
- Marshall, B. D. L. et al. (2016) 'Prevalence and Correlates of Lifetime Suicide Attempts Among Transgender Persons in Argentina', *Journal of Homosexuality*, 63(7), pp. 955–967. doi: 10.1080/00918369.2015.1117898.
- Mayer, K. H. et al. (2008) 'Sexual and gender minority health: what we know and what needs to be done.', *American Journal of Public Health*, 98(6), pp. 989–95.
- Meyer, I. H. (1995) 'Minority Stress and Mental Health in Gay Men', *Journal of Health and Social Behavior*, 36(1), p. 38.
- Meyer, I. H. (2003) 'Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence.', *Psychological Bulletin*, 129(5), pp. 674–697.
- Meyer, I. H. and Wilson, P. A. (2009) 'Sampling Lesbian, Gay, and Bisexual Populations', *Journal of Counseling Psychology*, 56(1), pp. 23–31.
- Müller, A. and Meer, T. (2018) *Access to justice for South African lesbian, gay, bisexual and transgender survivors of sexual assault*. Cape Town: GHJRU.
- Myer, L. et al. (2008) 'Common Mental Disorders among HIV-Infected Individuals in South Africa: Prevalence, Predictors, and Validation of Brief Psychiatric Rating Scales', *AIDS Patient Care and STDs*, 22(2), pp. 147–158.
- Nadal, K. L., Davidoff, K. C. and Fujii-Doe, W. (2014) 'Transgender Women and the Sex Work Industry: Roots in Systemic, Institutional, and Interpersonal Discrimination', *Journal of Trauma and Dissociation*, 15(2), pp. 169–183.
- Nemoto, T., Bödeker, B. and Iwamoto, M. (2011) 'Social support, exposure to violence and transphobia, and correlates of depression among male-to-female transgender women with a history of sex work', *American Journal of Public Health*, 101(10), pp. 1980–1988.
- OUT LGBT Well-being (2016) *Hate Crimes against Lesbian, Gay, Bisexual and Transgender (LGBT) people in South Africa*. Pretoria: OUT.
- Pega, F. and Veale, J. F. (2015) 'The Case for the World Health Organization's Commission on Social Determinants of Health to Address Gender Identity', *American Journal of Public Health*, 105(3), pp. e58–e62.
- Pinel, E. C. (1999) 'Stigma consciousness: The psychological legacy of social stereotypes.', *Journal of Personality and Social Psychology*, 76(1), pp. 114–128.
- Plöderl, M. and Tremblay, P. (2015) 'Mental health of sexual minorities. A systematic review'' *International Review of Psychiatry*, 27(5), pp. 367–385.
- Poteat, T. et al. (2017) 'HIV prevalence and behavioral and psychosocial factors among transgender women and cisgender men who have sex with men in 8 African countries: A cross-sectional analysis', *PLoS Medicine*, 14(11), pp. 1–17.
- Radloff, L. S. (1977) 'The CES-D Scale: A self report depression scale for research in the

- general population', *Applied Psychological Measurement*, 1(3), pp. 385–401.
- Robles, R. et al. (2016) 'Removing transgender identity from the classification of mental disorders: a Mexican field study for ICD-11', *The Lancet Psychiatry*, 3(9), pp. 850–859.
- Rosario, M. et al. (2014) 'Sexual-orientation disparities in substance use in emerging adults: a function of stress and attachment paradigms.', *Psychology of Addictive Behaviors : Journal of the Society of Psychologists in Addictive Behaviors*, 28(3), pp. 790–804.
- Rosser, B. R. S. et al. (2007) 'Capturing the social demographics of hidden sexual minorities: An internet study of the transgender population in the United States', *Sexuality Research and Social Policy*, 4(2), pp. 50–64.
- Sausa, L. A., Keatley, J. and Operario, D. (2007) 'Perceived risks and benefits of sex work among transgender women of color in San Francisco', *Archives of Sexual Behavior*, 36(6), pp. 768–777.
- Scheim, A. I., Bauer, G. R. and Shokoohi, M. (2016) 'Heavy episodic drinking among transgender persons: Disparities and predictors', *Drug and Alcohol Dependence*, 167, pp. 156–162.
- Sithole, B. (2017) 'HIV prevention needs for men who have sex with men in Swaziland', *African Journal of AIDS Research*, 16(4), pp. 315–320.
- Spitzer, R. L. et al. (2006) 'A Brief Measure for Assessing Generalized Anxiety Disorder', *Archives of Internal Medicine*. American Medical Association, 166(10), p. 1092.
- Stahlman, S. et al. (2015) 'Characteristics of men who have sex with men in Southern Africa who seek sex online: a cross-sectional study.', *Journal of Medical Internet Research*, 17(5), p. e129.
- Stevens, M. (2012) *Transgender access to sexual health services in South Africa: findings from a key informant survey*. Cape Town: Gender Dynamix.
- Suess Schwend, A. et al., 2018. Depathologising gender diversity in childhood in the process of ICD revision and reform. *Global Public Health*, 0(0), pp.1–14.
- Swaziland Government Ministry of Health (2014) 'WHO Steps Noncommunicable Disease risk Factor Surveillance Report: Swaziland 2014'. Available at: http://www.who.int/ncds/surveillance/steps/Swaziland_2014_STEPS_Report.pdf.
- The Other Foundation (2016) *Canaries in the Coal mines: An analysis of spaces for LGBTIQ activism in Southern Africa*, The Other Foundation. Available at: http://theotherfoundation.org/wp-content/uploads/2016/10/Canaries_Summary_epub_Draft4_MJ6.pdf.
- Victor, C. J. et al. (2014) 'The Psychological Society of South Africa sexual and gender diversity position statement: contributing towards a just society', *South African Journal of Psychology*, 44, pp. 292–302.
- Walters, M.L., Chen, J. & Breiding, M.J., 2013. The National Intimate Partner and Sexual Violence Survey 2010 Findings on Victimization by Sexual Orientation. Available at: https://www.cdc.gov/violenceprevention/pdf/nisvs_sofindings.pdf.
- World Health Organization (2003) *World Health Survey: Report of Swaziland*. Available at: <http://www.who.int/healthinfo/survey/whsswz-swaziland.pdf>.
- World Health Organization (2010) 'ATLAS of Substance Use Disorders. Resources for the Prevention and Treatment of Substance Use Disorders (SUD). Country Profile: Swaziland'.
- World Health Organization (2014a) 'Mental health Atlas country profile 2014: Swaziland'. Available at: http://www.who.int/mental_health/evidence/atlas/profiles-2014/swz.pdf?ua=1.
- World Health Organization (2014b) 'Swaziland alcohol consumption: levels and patterns (factsheet)'.
- World Health Organization (2017a) 'Depression and Other Common Mental Disorders: Global Health Estimates'. Geneva. Available at: <http://apps.who.int/iris/bitstream/handle/10665/254610/WHO-MSD-MER-2017.2-eng.pdf;jsessionid=0F7019C80E93A48089B95CCAD57B2DC5?sequence=1>.
- World Health Organization (2017b) 'Depression and Other Common Mental Disorders: Global Health Estimates'. Geneva: WHO.
- World Health Organization (2018a) *WHO ICD-11 Beta Draft*. Available at: <https://icd.who.int/dev11/l-m/en>.
- Young, R. M. and Meyer, I. H. (2005) 'The trouble with "MSM" and "WSW": erasure of the sexual-minority person in public health discourse.', *American Journal of Public Health*, 95(7), pp. 1144–9.

GLOSSARY OF TERMS RELATED TO SEXUAL ORIENTATION, GENDER IDENTITY AND EXPRESSION

Bisexual	People who are emotionally, romantically and/or sexually attracted not exclusively to people of one particular gender; attracted to both men and women.
Cisgender	Denoting or relating to a person whose sense of personal identity and gender corresponds with the sex assigned to them at birth.
Gay	A person who is emotionally, romantically and/or sexually attracted to persons of the same gender.
Gender expression	External appearance of one's gender identity, usually expressed through behaviour, clothing, haircut or voice, and which may or may not conform to socially defined behaviours and characteristics typically associated with being either masculine or feminine.
Gender identity	One's innermost concept of self as man, woman, a blend of both or neither – how individuals perceive themselves and what they call themselves. One's gender identity can be the same or different from their sex assigned at birth.
Gender minority	Gender minority refers to transgender and gender non-conforming/ gender diverse people whose gender identities or gender expressions fall outside of the social norms typically associated with the sex assigned to them at birth.
Gender non-conforming	A broad term referring to people who do not behave in a way that conforms to the traditional expectations of their gender, or whose gender expression does not fit neatly into a category.
Intersex	Intersex is an umbrella term for individuals who are born with sex characteristics that are, according to the typical understanding in society, either female and male at the same time, or not quite female or male, or neither female or male. This diversity can be related to chromosomes, hormones or anatomical features, and is not pathological.
Heterosexual	A person who is emotionally, romantically and/or sexually attracted to persons of the opposite gender.
Lesbian	Term used to describe female-identified people attracted romantically, sexually, and/or emotionally to other female-identified people.
LGBT, LGBTI	An acronym that refers to lesbian, gay, bisexual, transgender (and intersex if the 'I' is included) people. Often used together to refer to a shared marginalisation because of sexual orientation, gender identity and expression (and diversity of sex characteristics).
Sex assigned at birth	The assignment and classification of people as male, female, intersex, or another sex assigned at birth, often based on physical anatomy at birth and/or karyotyping.

Sexual activity	Sexual activity which includes sexual acts and sexual contacts, is the manner in which humans experience and express their sexuality.
Sexual attraction	Sexual attraction is attractiveness on the basis of sexual desire or the quality of arousing that interest. It is inherent to a person, and not a choice.
Sexual identity	Sexual identity is how someone thinks of him/herself in terms of to whom he/she is romantically or sexually attracted.
Sexual minority	A group whose sexual identity, orientation or practices differ from the majority of the surrounding society.
Sexual orientation	An enduring emotional, romantic, sexual or affectional attraction or non-attraction to other people. It is inherent to a person, and not a choice. Sexual orientation is not the same as gender identity.
Transgender	An umbrella term for people whose gender identity and/or expression is different from cultural expectations based on the sex they were assigned at birth. Being transgender does not imply any specific sexual orientation. Therefore, transgender people may identify as straight, gay, lesbian, bisexual, etc.
Transgender man	A person who identifies as a man, but was assigned a female sex at birth.
Transgender woman	A person who identifies as a woman, but was assigned a male sex at birth.

GLOSSARY OF TERMS RELATED TO THE STATISTICAL ANALYSIS

Adjusted Odds Ratio (AOR)	A statistical value that measures how strong an association between two variables might be. Odds ratio is a measure of association between an exposure and an outcome. Adjusted odds ratio is an Odds ratio which is adjusted for potential confounding by other variables.
Community-based sampling	Community-based sampling is a sampling methodology in which the researchers take their study participants (sample) from the community in general.
Confidence interval (CI)	Confidence intervals help us determine what the real value of a statistically calculated value might be. A confidence interval gives an estimated range of values which is likely to include an unknown population parameter, the estimated range being calculated from a given set of sample data.
Demographics	Properties of an individual or sample that can be regarded as factual, often used to structure a research sample. These include for example age, gender, sex, social class, working status and geographic location.
Descriptive statistics	Descriptive statistics are brief descriptive coefficients that summarize a given data set, which can be either a representation of the entire or a sample of a population. Descriptive statistics are broken down into measures of central tendency and measures of variability.
Electronic Data Management System (EDMS)	An Electronic Data Management System (EDMS) is a software package designed to manage electronic information and records within an organization's workflow.
Logistic regression model	Logistic regression is used to obtain odds ratio in the presence of more than one independent variable. It is used to analyse the relationship between two and more variables.
Mean	Mean is the most commonly used measure of central tendency. There are different types of mean inclusive of: arithmetic mean, weighted mean, geometric mean, and harmonic mean. If mentioned without an adjective (as mean), it generally refers to the arithmetic mean, which is computed by adding all the values in the data set divided by the number of observations in it.
Multiple imputation	Multiple imputation is a general approach to the problem of missing data that is available in several commonly used statistical packages. It aims to allow for the uncertainty about the missing data by creating several different plausible imputed data sets and appropriately combining results obtained from each of them.
Online-based sampling	Online-based sampling is a sampling method from a population of individuals when the primary method of gathering the responses to a given survey comprising a set of questions contained in a questionnaire with the purpose of identifying the attitudes of the given population, is over the Internet.

p-value	The p-value or probability value is a statistical test to assess if what we can see in the data is there by chance. The smaller the p value, the less likely it is that what we see in the data is coincidental.	
Pilot survey	A pilot survey is conducted with few individuals of the target population or the sample of a survey, in order to test and refine the survey instruments (questionnaire and instruction manual, data processing manual and programmes) before the main data collection starts across the target population or the full sample.	
Prevalence	Prevalence refers to the total number of individuals in a population who have a disease or health condition at a specific period of time, usually expressed as a percentage of the population.	
Protocol	A (research) protocol is a detailed document that describes the background, rationale, objectives, design, methodology, statistical considerations, and organization of a clinical research project.	
Protocol violation	A divergence from the protocol that reduces the quality or completeness of the data, makes the Informed Consent Form inaccurate, or impacts a participant's safety, rights, or welfare.	
Sample	In statistics, a sample refers to a set of observations drawn from a population.	
Sample size	Sample size is the number of observations in a sample, often denoted with "n". It describes the number of participants who have filled out a survey, and whose answers have been taken into account when analysing the data.	
Survey	A survey is an investigation about the characteristics of a given population by means of collecting data from a sample of that population and estimating their characteristics through the systematic use of statistical methodology.	
Questionnaire administration	The process of asking questions and recording the answers.	
	Self-administration	When the questionnaires are read and filled by the respondents themselves, the questionnaire administration is called self-administration.
	Fieldworker-administration	When a fieldworker read the questions to the participant, the questionnaire administration is called Fieldworker-administration.
Variable	A variable is a characteristic of a unit being observed which may assume more than one of a set of values, to which a numerical measure or a category from a classification can be assigned.	
	Binary variable	A binary variable is a variable with only two values.
	Continuous variable	A continuous variable is a variable that has an infinite number of possible values.

APPENDIX 1: DETAILED METHODOLOGY

Measures: Sexual orientation and gender identity

Survey questions

In order to paint a nuanced picture of the participants' sexual orientation, we aimed to assess self-identified sexual identity, sexual attraction and sexual behaviour. We asked the following questions:

4. **Self-identified sexual identity** was assessed by asking participants "In terms of your sexual orientation, how do you identify?" (Options: Lesbian, Bisexual, Gay, Heterosexual, Asexual, "Other, specify")
5. **Attraction** was assessed by asking participants who they were sexually and emotionally attracted to (2 questions).
6. **Sexual activity** was assessed by asking participants about who they have had "sexual experiences with in the past year and their lifetime" (2 questions).

For attraction and sexual activity, the questionnaire gave participants a list of options from which they could select all that applied (Options: With women, with men, with trans women, with trans men, with gender non-conforming people, with intersex people, "I have not had sexual experiences", "Other, specify").

To measure a participant's gender identity, we combined three questions:

1. **Self-identified gender identity** was assessed by asking "In terms of your gender identity, how do you identify?" (Options: Woman, Man, Trans woman, Trans man, Gender non-conforming, "Other, specify").
2. We asked about **sex assigned at birth** (Options: Male, Female, Intersex)
3. Additionally, we asked what sex/ gender was recorded in the participant's identity document(s)

Categorisation for analysis

Throughout this report, we use categories of sexual orientation (lesbian, gay, bisexual, 'non-normative', and heterosexual) and gender identity (cisgender women, cisgender men, transgender women, transgender men and gender non-conforming people) to disaggregate the findings about experiences of violence and mental health outcomes. To create these categories, we in some instances had to re-code the way participants self-identified, based on the other information they provided in the questions about their sexuality and gender identity. Re-coding in these categories was done in the following ways:

Sexual orientation

- *Lesbian (and other women who have sex with women)*: any participant who identified 'lesbian' as their sexual orientation; any cisgender woman who identified 'gay' as their sexual orientation; any transgender woman who identified as 'gay' and was sexually attracted to/has sex with women; any transgender man who identified as 'gay' and was sexually attracted to/has sex with women⁷; any cisgender or transgender woman who identified as 'heterosexual' but exclusively had sex with women in the past year; any cisgender or transgender woman who identified as 'heterosexual,' had not had sex with anyone in the past year and was exclusively sexually attracted to women; gender non-conforming people who identify as gay and have sex exclusively with women.
- *Gay (and other men who have sex with men)*: Any transgender or cisgender man, gender non-conforming person, or 'other' gender identity who identified their sexual orientation as 'gay'; any transgender woman who identified as 'gay' and was sexually attracted to/has sex with men⁸; men who identified their sexual orientation as 'homosexual' or 'MSM'; any cisgender or transgender man who identified as 'heterosexual' but exclusively had sex with men in the past year; any cisgender or transgender man who identified as 'heterosexual,' had not had sex with anyone in the past year and was exclusively sexually attracted to men.
- *Bisexual*: any participant who identified as 'bisexual'.
- *Non-normative sexual orientation*: We were cognisant that the more widely used sexual orientations (lesbian, gay, bisexual) depend on the assumption of a gender binary: one can only classify their sexual orientation if one's own gender and one's partner's gender is either woman or man; ie. lesbian means that one identifies as a woman and is attracted to or has sex with other women (Better and Simula, 2015). If one's partner identifies as gender non-conforming, it is not possible to classify one's sexual orientation as lesbian (a woman attracted to women), gay (a man attracted to men) or bisexual (a woman or a man attracted to both men and women). For those participants whose sexual orientation transgressed the gender binary, and for participants who did not fit the gender binary needed to classify their sexual orientation as lesbian, gay or bisexual, we created a new category: that of 'non-normative' sex orientation. The 'non-normative' indicates that they could not be classified as any of the more widely used sexual orientations (lesbian, gay or bisexual). A lot of these participants had listed their sexual orientation as 'other' – including for example, queer or pansexual. Additionally, it includes participants who identified as 'heterosexual' and who reported having sex with people of more than one sex/gender in the past year.
- *Heterosexual*: any participant who identified as 'heterosexual' and had sex with only people of a different sex/gender in the past year.

Gender identity

- *Transgender women*: Those who self-identified as trans women; those who self-identified

7 Transgender men who had sex with women and identified as heterosexual were grouped as 'heterosexual'. While grouping transgender men who identify as gay and who are attracted to and have sex with women as 'lesbian' does not completely accurately capture their self-defined identity, we felt it would have been even less accurate to group them with cisgender men who have sex with men.

8 See previous footnote. Transgender women who had sex with men and identified as heterosexual were grouped as 'heterosexual'. While grouping transgender women who identify as gay and who are attracted to and have sex with men as 'gay' does not completely accurately capture their self-defined identity, we felt it would have been even less accurate to group them with cisgender women who have sex with women.

- as women and were assigned male at birth.
4. *Transgender men*: those who self-identified as trans men; those who self-identified as men and were assigned female at birth.
 5. *Gender non-conforming*: those who self-identified as gender non-conforming, regardless of sex assigned at birth.

Measures: Mental health

CES-D 10: Depression

We used the instrument CES-D 10, a 10-item Center for the Epidemiological Studies of Depression Short Form to measure depression. It is widely used to screen for signs of depression in primary care settings, and is often used for research on the prevalence of depression. It is important to keep in mind, however, that we cannot diagnose people using the CES-D 10. In order to receive a definitive diagnosis of clinical depression, an individual needs to see a healthcare provider.

We followed the CES-D 10 instructions to categorise scores into a binary variable, using a cut-off score of 10, where participants with a CES-D 10 score of 10 or above were considered to have signs of depression and those with a score under 10 were classified as not having signs of depression. Additionally, we report only on participants who had no more than two missing values on the CES-D 10 items (Radloff, 1977). However, for logistic regression models including CES-D 10 as a covariate, the continuous variable of the CES-D 10 score was used and multiple imputation was used for missing values. For the logistic regression model where the CES-D 10 score was the outcome, the binary variable was used.

GAD-7: Anxiety

The Generalized Anxiety Disorder 7-item scale (GAD-7) uses seven scored Likert items that assess signs of anxiety in the last two weeks. We created a categorical variable with the following cut-off scores: score of 0 to 4 indicates no anxiety symptoms; score of 5 to 9 indicates mild anxiety symptoms; score of 10 to 14 indicates moderate anxiety symptoms; score of 15 or above indicates severe anxiety symptoms. We also created a binary variable using a score of 10 as a cut-off to compare no/mild anxiety with moderate/severe anxiety, which was used for the logistic regression model where GAD-7 score was the outcome (Kroenke, Spitzer and Williams, 2001; Spitzer *et al.*, 2006). We excluded participants who had missing data for any GAD-7 items from GAD-7 scoring. In logistic regression models in which GAD-7 was a covariate, we used the continuous GAD-7 score, and used multiple imputation to impute missing data.

AUDIT: Alcohol

The Alcohol Use Disorders Identification Test (AUDIT) uses 10 items to assess whether an individual's alcohol use is harmful. The questions ask about how often participants drink alcohol, how much, and how their alcohol use has impacted their life (e.g. "Have you or someone else been injured because of your drinking?"). Participants who do not drink have an AUDIT score of 0. For those who do drink, we followed the AUDIT manual to create a categorical variable with the following cut-offs: score of 1 to 7 indicates non-hazardous alcohol use; score of 8 to 15

indicates hazardous use; score of 16 to 19 indicates harmful use; score of 20 and above indicates alcohol dependence. We excluded participants who had missing data for any AUDIT items from AUDIT scoring. For the logistic regression model where AUDIT was the outcome, we used a binary variable with a cut-off score of 8 (Barbor *et al.*, 2001). In logistic regression models in which AUDIT was a covariate, we used the continuous AUDIT score. We used multiple imputation to impute missing data for the regression models.

DUDIT: Drugs

The Drug Use Disorders Identification Test (DUDIT) is a scale with 11 items to assess harmful drug use. We created a categorical variable using the following categories, which are suggested by the DUDIT manual: score of 0 for those who do not do drugs; score of 1 to 5 for some drug use; score of 6 to 24 for harmful use; score of 25 and above indicates drug dependence (on one or more drugs) (Berman *et al.*, 2003). To create a binary variable, the DUDIT manual recommends different cut-off scores for men and women, and does not specify what to do in instances of gender minority people. Recognising the limitations of these recommendations for a study with gender diverse participants, we chose to use the higher cut-off score of 6, which the manual recommends for men, for participants of all genders. We used the binary variable with this cut-off point in the logistic regression model where DUDIT was the outcome. In logistic regression models in which DUDIT was a covariate, we used the continuous DUDIT score. We excluded participants who had missing data for any DUDIT items from DUDIT scoring, however we used multiple imputation to impute missing data in the regression models.

Signs of post-traumatic stress

We created a binary variable for signs of post-traumatic stress: those who said they experienced all three signs were categorised as having signs of post-traumatic stress; those who said they experienced one, two, or no signs were categorised as not having signs of post-traumatic stress. This binary variable was used when post-traumatic stress was included as a co-variate in logistic regression models.

Sampling and enrolment

Decisions around sampling for LGBTI populations are complex, and impacted by a number of factors unique to this population and the specific country-context. Sampling is complicated by the following factors, as described by Meyer and Wilson (Meyer and Wilson, 2009):

- LGBTI populations are not easy to identify. Sexual orientation and gender identity are not fixed constructs, different people have different identities, and this is particularly important in contexts where Western concepts of L, G, B, T and I might not hold the same value for everybody. Further, many LGBTI people may not reveal their gender or sexual orientation, or seek assistance from LGBTI organisations, for fear of discrimination.
- LGBTI populations are hidden. For a sampling method that predicts larger, population-size trends, researchers need to know the overall population size, in our example, the overall number of LGBTI individuals in each country. This of course is impossible to determine, both because of the previous point, and because sexual orientation and gender identity are not registered in national census data, thus making it impossible to obtain this information.

This means that sampling methods that will allow us to make predictions about ALL LGBTI people in a certain context are impossible at this moment.

- Given that many partner organisations do not have definite numbers of their constituency population, it would be impossible for us to even make generalising predictions about any organisations’ constituency population, for the same reasons outlined in the previous point (Meyer & Wilson, 2009).

Given these restrictions, we combined two sampling methods: community-based sampling and online-based sampling. We chose to combine these two sampling methods for two reasons:

- Hendricks and Testa (Hendricks and Testa, 2012) show that needs assessments and community-based samples, such as the one we used for our study, often reach especially vulnerable parts of sexual and gender minority populations. This means that the people who participate in community-based surveys, such as ours, are often disadvantaged in more than one way, and so face oppression on more than one level. This means that what we learn from community-based sampled studies can illustrate minority stress by reaching those who are most affected.
- However, Rosser and colleagues (Rosser et al., 2007) have pointed out the limitations of community sampling, which may over-represent targeted problems. In our sample, this means that by sampling people who already access NGOs (arguably because they feel they need support), we might over-estimate the level of mental health problems among sexual and gender minority people more generally. Therefore, we have added online-based sampling to also reach people who do not access NGO services directly.⁹

The following table provides an overview of the number of participants in each country, as well as the number of participants enrolled by each organisation.

Partner organisation	Number of participants
Botswana	618
Bonela	223
LeGaBiBo	168
RIA	221
Other (filled out in Kenya but living in Botswana)	3
Ethiopia	198
Organisation 1	64
Organisation 2	119
Other (online)	15

⁹ In some countries, the online response rate was poor, or partner organisations chose not to implement online data collection. This was for various reasons, including: poor access to internet, poor access to data collection devices and safety concerns about publicising a public survey link. We describe the country-specific use of the online survey in the Findings section.

Partner organisation	Number of participants
Kenya	976
Ishtar-MSM	183
Jinsiangu	76
Maaygo	181
Minority Women in Action	104
National Gay and Lesbian Human Rights Commission	215
PEMA	216
Other (online)	1
Lesotho	173
People's Matrix Association	173
Malawi	197
Centre for the Development of the People	196
Other (collected in Kenya, participant living in Malawi)	1
South Africa	832
Durban Lesbian and Gay Community and Health Centre	102
Gender Dynamix	166
OUT LGBT Well-Being	202
Triangle Project	256
Other (online)	106
eSwatini	103
Rock of Hope	102
Other (online)	1
Zambia	353
Friends of Rainka	197
TransBantu Zambia	59
The Lotus Identity	90
Other (online)	7

Partner organisation	Number of participants
Zimbabwe	346
Gays and Lesbians of Zimbabwe	178
Sexual Rights Centre	165
Other (online)	3
TOTAL	3,796

Data management

Once the partner organisations had finished collecting data, all questionnaires were sent to the GHJRU's offices at the University of Cape Town for data entry. Data were entered by trained research assistants, using the RedCap online survey tool.

Data quality

We undertook a number of steps to ensure that the quality of data was as high as possible. Questionnaires with good data quality are questionnaires that are completely filled out.

For the online survey: The REDCap online survey had checks for data quality in place. For example, skip/logic patterns were programmed into the survey. The online survey also prompted participants to fill out questions that they had accidentally left out.

For the paper survey: We trained fieldworkers to review all completed paper surveys before the participant who had filled it out left. This was so that the fieldworker could identify questions that the participant might have missed, or questions that the participant should not have answered, or questions where the participant had ticked more than one answer. Because the survey was totally anonymous, we could not go back to participants and ask them about questions they had not filled out, or questions that they had filled out incorrectly (where, for example, they had ticked two possible answers and we did not know which one was correct).

Once received at the GHJRU offices, we (the researchers) checked all surveys checked for quality. We trained people to enter the data, who would also identify unusual responses or errors in the data documented on the surveys. When necessary, we held meetings with the data enterer to decide on "data entry rules" for surveys where participants had ticked contradictory answers. We applied these data entry rules to all surveys.

In cases where the participants had not ticked yes to all eligibility questions, or where they had not ticked yes to say that they consented to participating, we did not enter the data from the survey and excluded the participant from the study.

Data cleaning

We used REDCap was used during the data cleaning process to update data in instances of data entry error. Following this, data was exported to Stata. We used Stata to examine patterns of missing and conflicting data. Unusual or unexpected responses that were identified in this process were checked against paper copies and amended as needed.

“Other, specify” responses were reviewed by the research team. We recorded decisions on how to code these write-in responses in the “data entry rules,” which were applied to data from all countries. In instances of large numbers of the same “other” responses, we created new coding categories.

Conflicting data

In some instances, questions asked about the same experience twice: first about the experience in participants’ lifetime, then in the last 12 months. For example:

301.	Has there ever been a period of time when you thought about committing suicide?	In your lifetime?	₁ Yes	₀ No
		In the last 12 months?	₁ Yes	₀ No

In some instances, participants entered a conflicting response; for example, saying that they had not thought about suicide in their lifetime, but had thought about it in the last 12 months. In some instances, they left the question about lifetime incomplete, but said they had thought about suicide in the last 12 months. During data cleaning, we made the decision to recode “lifetime” as “yes” in both these instances – so if a participant said they had experienced something in the past 12 months, by default they had also experienced it in their lifetime. This was done for all questions in the above format in the questionnaire.

Data analysis

All data from the online survey and paper survey were managed through REDCap at the University of Cape Town. Data cleaning was completed with REDCap and Stata15. Data analysis was conducted with Stata15.

Describing the data

The main aim of this research was to report prevalence of mental health concerns, healthcare access experiences, experiences of violence, social support and stigma among sexual and gender minority people in our sample.

For this reason, the majority of the report uses descriptive statistics to explain what the research participants reported. These findings should not be considered “representative” of the sexual and gender minority population in each country. However, as an exploratory, cross-sectional study we hope that our findings will reveal priority areas for future research and service delivery, considering the dearth of evidence on sexual and gender minority people’s mental health and wellness on the continent.

Measuring associations

This study did not collect information from heterosexual, cisgender people. Because of this, our findings do not report on sexual and gender minority people as compared to their heterosexual, cisgender counterparts. In some instances we drew on peer-reviewed and grey literature in order to discuss our findings as compared to other populations.

In some instances, we report on interesting associations we found within our own sample. For example, we often examined differences between gender minorities and cisgender participants (where the cisgender participants are sexual minority people) and between black and white participants (where black refers to any participant who did not identify as white). For these comparisons, we started with using chi squared (or Fisher's exact) tests to assess raw associations between categories. The p-values for these tests are reported in tables throughout the Findings section of this report. P-values describe the statistical significance of the association, that is, the chances of whether the association we found is simply due to chance.

Logistic regression

In some instances, we used a tool called logistic regression to examine differences in outcomes within our sample. For example, in countries with large sample sizes, we used logistic regression to assess if there was a difference in depression level ('outcome') between cisgender and gender minority participants ('predictor') while also accounting for other factors.

Logistic regression is used when an outcome has multiple predictors (factors that may cause, prevent or contribute to the outcome). By using logistic regression, we are able to measure association between the outcome and multiple predictors at the same time. Logistic regression produces adjusted odds ratios (AORs), which measures the size of association between different predictors and the outcome.

In our logistic regression models, we included predictors that are known or suspected confounders ("third variables" that influence both a predictor and an outcome) or that are believed to otherwise influence the outcome. This inclusion is called 'adjustment', meaning that the AOR takes into account the effects of other predictors when describing the relationship between any one predictor and outcome.

Examining the AOR gives information about how predictors and outcomes were related in our sample. AORs greater than 1 mean that as the predictor increases, the odds of the outcome increases ("positively associated") and AORs less than 1 mean that as the predictor increases, the odds of the outcome decreases ("negatively associated").

P-values and confidence intervals add understanding about whether these findings are due to chance. A p-value is a measure related to probability. The confidence interval expresses a range in which we are "confident" that the true AOR exists. For this study, we used 95% confidence intervals for AORs—meaning that we are 95% confident that the 'true' association between the predictor and outcome lies within the confidence interval. A p-value of less than 0.05 indicates that there is a 'true' difference in the outcome as a predictor changes (while also accounting for the other predictors in the model).

Example

For example, in South Africa, we found that lifetime experience of sexual violence was associated with suicidal ideation in the last year (see in the South Africa section of this report):

Suicidal ideation (last year)	AOR	95% CI	p
No experience of sexual violence	-	Reference category	
Experienced sexual violence (lifetime)	2.05	1.29 – 3.26	0.003

We can interpret this table as follows:

- Reference category is “no experience of sexual violence” – this means that the predictor is “experienced sexual violence (lifetime)”, which will be compared to “no experience of sexual violence” (the reference category)
- AOR of 2.05 – The odds of suicidal ideation in the last year are 2.05 greater in those who experienced lifetime sexual violence, in comparison to those who did not experience sexual violence, holding all other factors constant.
- 95% confidence interval of 1.29-3.26 – We are 95% confident that the AOR is between 1.29 and 3.26.
- p-value of 0.003 – The p-value is less than 0.05 (<0.05) which means we believe that there is a statistically significant difference in the AOR of suicidal ideation in the last year between those who have and have not experienced sexual violence in their lifetimes.

Missing data

Prior to beginning analysis, we examined patterns of missing data. Missing data was sometimes more common for specific variables than others.

Due to the anonymous nature of the questionnaire, we could not follow-up with participants to ask their response when a questionnaire item was incomplete. We recorded these in the database as missing data.

Missing data was more common in the “outcomes” section of the questionnaire, which came after demographics, and among those who completed the questionnaire online. We expect that some participants chose to end the survey early or were otherwise interrupted while completing the online survey. In analysis, we included only questionnaires (paper and online) in which the participant completed at least some items in the “outcomes” section.

Patterns of missing data were different between study countries, study sites, and between questionnaire items. After consideration, we decided to report descriptive statistics using only complete data (please note the sample sizes in the “Findings” of this report by locating the “n” for each table or figure). This is known as “complete case analysis.”

For some measures of association, we utilised a method for dealing with missing data called multiple imputation. Multiple imputation is a statistical process with three steps: (1) imputation—statistical software is used to generate duplicate datasets in which the missing data has been replaced by calculated values (“imputations”), (2) analysis—each imputed data set is analysed separately, (3) pooling—the separate analyses are statistically pooled into one measure of association.

Multiple imputation is useful because it can help prevent bias that missing data can cause.

We decided not to apply multiple imputation while reporting on descriptive statistics, although this has been done by others elsewhere. Based on the designed purpose of multiple imputation, imputed data is not meant to truly replace or substitute the answer that would have been true for a participant. Rather, imputed data is used more like a place holder so that a statistical analysis can be stronger. For this reason, we felt that reporting imputed data in descriptive statistics would be misleading.

We used multiple imputation to account for missing data in all regression models. To multiply impute, we used predictive mean matching for continuous variables and categorical scale items (i.e. Likert scales) and logistic regression for binary variables. Predictive mean matching was a method designed for continuous data, but it has been suggested it can also be applied to categorical variables (Morris, White and Royston, 2014). We imputed only variables that were necessary for these analyses, as well as additional variables we felt might be associated with “missingness” of data. All variables relevant to the analyses were imputed, even when the amount of missing data was small.

APPENDIX 2: QUESTIONNAIRE

SWAZILAND—ENGLISH

Instructions for self-administration

You will complete this questionnaire by yourself. A fieldworker will review what the study is about and check that you are eligible and willing to be in the study.

Carefully complete this questionnaire. Check that you have completed every question.

For most questions, choose one response.

106.	Do you own your housing? PLEASE TICK ONE	<input type="radio"/> 1 Yes, I own it myself <input type="radio"/> 2 No, I rent it <input checked="" type="radio"/> 3 No, I share housing and do not pay for it <input type="radio"/> 4 Not applicable (living on the street)			
208.	When seeking healthcare, how often do you think you have been treated disrespectfully by staff for being LGBTI? (this includes doctors, nurses, counselors, other people working at public, private, or traditional healthcare facilities)	<input checked="" type="radio"/> 1 Never	<input type="radio"/> 2 Rarely	<input type="radio"/> 3 Sometimes	<input type="radio"/> 4 Often
209.	When seeking healthcare, how often do you think have you received poorer service than other people for being LGBTI?	<input type="radio"/> 1 Never	<input checked="" type="radio"/> 2 Rarely	<input type="radio"/> 3 Sometimes	<input type="radio"/> 4 Often

Some items allow you to tick more than one response.

112.	Who do you feel sexually attracted to? PLEASE TICK ALL THAT APPLY	<input checked="" type="checkbox"/> 1 To women <input type="checkbox"/> 2 To men <input type="checkbox"/> 3 To trans women <input checked="" type="checkbox"/> 4 To trans men <input type="checkbox"/> 5 To gender non-conforming people <input checked="" type="checkbox"/> 6 To intersex people <input type="checkbox"/> 7 I do not feel sexual attraction <input type="checkbox"/> 8 Other, specify: _____				
------	--	--	--	--	--	--

Sometimes the same question is asked twice—once about the last 12 months and once about your whole lifetime (ever).

403.	Has anyone ever <u>insulted or verbally harassed</u> you because of being LGBTI?	a. In your life time?	<input checked="" type="radio"/> 1 Yes	<input type="radio"/> 2 No
		b. In the last 12 months?	<input type="radio"/> 1 Yes	<input checked="" type="radio"/> 2 No

Be sure to answer **both** questions. Remember that if you experienced something in the last 12 months, you have also experienced it in your lifetime.

If you make a mistake, make the correction clearly. Place one or two lines through the incorrect response and circle the correct response.

214.	Have you postponed or not tried to get needed healthcare when you were sick or injured because you could not afford it?	<input checked="" type="radio"/> 1 Yes	<input type="radio"/> 2 No
215.	Have you postponed or not tried to get needed healthcare because you could not afford it?	<input checked="" type="radio"/> 1 Yes	<input type="radio"/> 2 No

SWAZILAND—ENGLISH

Questionnaire consent statement

The Gender Health and Justice Research Unit at the University of Cape Town, in partnership with COC Netherlands and community based organisations across 12 African countries, (Angola, Botswana, Kenya, Lesotho, Malawi, Mozambique, Namibia, South Africa, Swaziland, Tanzania, Zambia, and Zimbabwe) is surveying people who are/identify as LGBTI. We aim to develop a better understanding of mental health, wellbeing, and experiences of discrimination, in order to inform advocacy efforts for improved services.

For this survey, we use LGBTI to mean someone who is or identifies as *any* of the following: gay, lesbian, bisexual, pansexual, omnisexual, asexual, men who have sex with men (MSM), women who have sex with women (WSW), transgender, transsexual, transman, transwoman, non-binary trans, queer, genderqueer, gender diverse, gender non-conforming, intersex and body diverse.

Please fill out our quick survey to let us know about your experiences accessing healthcare, about your mental health and well-being, and your experiences with violence.

This survey should take about 20-30 minutes to complete. This survey is **anonymous**, meaning that we will not ask for your name or any other identifying information. What you share in this survey will be kept confidential.

At the end of this survey, we will include a list of resources in your country should you need someone to talk to about your mental health, wellbeing, or experiences of discrimination.

The outcomes of the survey will be used to inform agenda setting by the COC Netherlands and in-country partner organisations to plan advocacy efforts around improving access to services for LGBTI people, particularly mental health services. The findings of this study may be published in academic literature, in which case your answers will not be linked to any identifying information. We can email you a report with the outcomes of this survey. If you wish to receive this report, please contact the organisation that gave you this questionnaire or sent you the link.

Please do not hesitate to contact us if you have any questions. If you have questions about your rights as a research participant, please contact the Faculty of Health Sciences Human Research Ethics Committee, Room E52-54 Groote Schuur Hospital Old Main Building, Observatory 7925, phone +27 21 406 6338 or email shuretta.thomas@uct.ac.za.

To begin, please complete the eligibility questions below.

Thank you for your assistance.

Kind regards

Dr Alex Muller
Senior Researcher
Gender Health and Justice Research Unit
University of Cape Town
Falmouth Building, Entrance 1, Level 1, Room 1.01.5
(021) 406 6021
alexandra.muller@uct.ac.za

These questions should be completed by a fieldworker:

1. Are you 18 years of age or older?

- 1 Yes
- 0 No → NOT ELIGIBLE

2. Do you identify as LGBTI (see above)?

- 1 Yes
- 0 No → NOT ELIGIBLE

3. Do you currently live in Angola, Botswana, Kenya, Lesotho, Malawi, Mozambique, Namibia, South Africa, Swaziland, Tanzania, Zambia, or Zimbabwe?

- 1 Yes
- 0 No → NOT ELIGIBLE

This question should be *ticked by the participant*, but can be asked by a fieldworker:

4. Do you agree to participate in this survey, based on the information outlined above? (this will be regarded as your informed consent to participate in this survey)

- 1 Yes
- 0 No → NOT ELIGIBLE

5. Are you completing the questionnaire by yourself?

- 1 Yes (self-administered)
- 0 No (fieldworker administered)

The following question should be completed by the fieldworker.

6. Has the participant answered yes to questions 1, 2, 3 and 4?

- 0 No → Sign and STOP HERE. Explain to participant they are not eligible for the survey. Place this completed form in a secure place.
- 1 Yes → Sign and continue data collection per guidelines in the Fieldworker Manual.

Fieldworker signature: _____ Date: _____

SWAZILAND—ENGLISH

Section 1a: Background

101.	How old are you?	PLEASE WRITE YOUR AGE: _____
102.	In which country do you currently live? PLEASE TICK ONE	<input type="radio"/> 1 Angola <input type="radio"/> 2 Botswana <input type="radio"/> 3 Kenya <input type="radio"/> 4 Lesotho <input type="radio"/> 5 Malawi <input type="radio"/> 6 Mozambique <input type="radio"/> 7 Namibia <input type="radio"/> 8 South Africa <input type="radio"/> 9 Swaziland <input type="radio"/> 10 Tanzania <input type="radio"/> 11 Zambia <input type="radio"/> 12 Zimbabwe
103.	How did you hear about this study?	<input type="radio"/> 18 Swaziland for Positive Living (SWAPOL) <input type="radio"/> 19 The Rock of Hope (ROH) <input type="radio"/> 20 House of Our Pride
104.	How do you identify your race?	<input type="radio"/> 1 Black <input type="radio"/> 2 White <input type="radio"/> 5 Other specify: _____
105.	In what type of housing do you currently live?	<input type="radio"/> 1 House <input type="radio"/> 2 Apartment / flat <input type="radio"/> 7 Room <input type="radio"/> 3 Shanty / Shack <input type="radio"/> 4 Hotel <input type="radio"/> 5 Mobile house <input type="radio"/> 6 On the street <input type="radio"/> 8 Hut/Traditional dwelling

SWAZILAND—ENGLISH

106.	Do you own your housing? PLEASE TICK ONE	<input type="radio"/> 1 Yes, I own it myself <input type="radio"/> 2 No, I rent it <input type="radio"/> 3 No, I share housing and do not pay for it <input type="radio"/> 7 Not applicable (living on the street)
107.	What type of area do you live in?	<input type="radio"/> 1 Urban <input type="radio"/> 2 Semi-urban/Peri-urban <input type="radio"/> 3 Rural
108.	On average, do you have enough money to cover your basic needs?	<input type="radio"/> 1 Yes <input type="radio"/> 0 No
109.	Do you have a job for which you are paid?	<input type="radio"/> 1 Yes, I have formal employment (I have an employment contract) <input type="radio"/> 2 Yes, I have informal employment (I am paid for work but do not have an employment contract) <input type="radio"/> 0 No, I do not have any work for which I am paid
110.	Which religion, if any, most closely aligns to your beliefs?	<input type="radio"/> 1 African tradition <input type="radio"/> 2 Islam <input type="radio"/> 3 Christianity <input type="radio"/> 4 Rastafarianism <input type="radio"/> 5 Judaism <input type="radio"/> 6 I am not religious <input type="radio"/> 7 Other, specify: _____
111.	What is the highest level of education that you have completed?	<input type="radio"/> 1 No formal education <input type="radio"/> 2 Primary education <input type="radio"/> 3 Secondary school <input type="radio"/> 4 Post-secondary school/University diploma or degree

SWAZILAND—ENGLISH

112.	Who do you feel sexually attracted to? PLEASE TICK <u>ALL</u> THAT APPLY	<input type="checkbox"/> 1 To women <input type="checkbox"/> 2 To men <input type="checkbox"/> 3 To trans women <input type="checkbox"/> 4 To trans men <input type="checkbox"/> 5 To gender non-conforming people <input type="checkbox"/> 6 To intersex people <input type="checkbox"/> 7 I do not feel sexual attraction <input type="checkbox"/> 8 Other, specify: _____
113.	Who do you feel emotionally attracted to? PLEASE TICK <u>ALL</u> THAT APPLY	<input type="checkbox"/> 1 To women <input type="checkbox"/> 2 To men <input type="checkbox"/> 3 To trans women <input type="checkbox"/> 4 To trans men <input type="checkbox"/> 5 To gender non-conforming people <input type="checkbox"/> 6 To intersex people <input type="checkbox"/> 7 I do not feel emotional attraction <input type="checkbox"/> 8 Other, specify: _____
114.	In the last year, whom have you had sexual experiences with? PLEASE TICK <u>ALL</u> THAT APPLY	<input type="checkbox"/> 1 With women <input type="checkbox"/> 2 With men <input type="checkbox"/> 3 With trans women <input type="checkbox"/> 4 With trans men <input type="checkbox"/> 5 With gender non-conforming people <input type="checkbox"/> 6 With intersex people <input type="checkbox"/> 7 I have not had sexual experiences in the last year <input type="checkbox"/> 8 Other, specify: _____
115.	In your lifetime, whom have you had sexual experiences with? PLEASE TICK <u>ALL</u> THAT APPLY	<input type="checkbox"/> 1 With women <input type="checkbox"/> 2 With men <input type="checkbox"/> 3 With trans women <input type="checkbox"/> 4 With trans men <input type="checkbox"/> 5 With gender non-conforming people <input type="checkbox"/> 6 With intersex people <input type="checkbox"/> 7 I have never had sexual experiences <input type="checkbox"/> 8 Other, specify: _____

SWAZILAND—ENGLISH

116.	<p>In terms of your sexual orientation, how do you identify?</p> <p>PLEASE TICK ONE</p>	<p><input type="radio"/> 1 Lesbian</p> <p><input type="radio"/> 2 Bisexual</p> <p><input type="radio"/> 3 Gay</p> <p><input type="radio"/> 4 Heterosexual</p> <p><input type="radio"/> 5 Asexual</p> <p><input type="radio"/> 6 Other; please specify _____</p>
117.	<p>In terms of your gender identity, how do you identify?</p> <p>PLEASE TICK ONE</p>	<p><input type="radio"/> 1 Woman</p> <p><input type="radio"/> 2 Man</p> <p><input type="radio"/> 3 Trans woman</p> <p><input type="radio"/> 4 Trans man</p> <p><input type="radio"/> 5 Gender non-conforming</p> <p><input type="radio"/> 6 Other; please specify: _____</p>
118.	<p>How was your sex classified at birth?</p> <p>PLEASE TICK ONE</p>	<p><input type="radio"/> 1 Female</p> <p><input type="radio"/> 2 Male</p> <p><input type="radio"/> 3 Intersex (persons born with sex organs/genitals that do not appear typically female or typically male)</p>
119.	<p>What is the legal sex/gender currently recorded in your identity document?</p> <p>PLEASE TICK ONE</p>	<p><input type="radio"/> 1 Female</p> <p><input type="radio"/> 2 Male</p> <p><input type="radio"/> 3 Intersex</p> <p><input type="radio"/> 4 Unspecified</p> <p><input type="radio"/> 5 Other; please specify: _____</p> <p><input type="radio"/> 77 I do not have an identity document</p>

SWAZILAND—ENGLISH

Section 1b: Gender expression

We would now like to know more about your gender expression. Indicate on a scale from 1 (not at all) to 5 (extremely) how masculine and feminine *you think you are*. We understand that being masculine or feminine is not natural or something you are born with, but we would like to know about how much you conform to society's expectations of what is masculine or feminine.

Place an X in one box that best describes your answer to each question.

120.	In general, how feminine do you think you are?	₁ Not at all	₂ A little	₃ Somewhat	₄ Very much	₅ Extremely
121.	In general, how feminine do you behave in front of others?	₁ Not at all	₂ A little	₃ Somewhat	₄ Very much	₅ Extremely
122.	In general, how feminine do you appear to others?	₁ Not at all	₂ A little	₃ Somewhat	₄ Very much	₅ Extremely
123.	In general, how masculine do you think you are?	₁ Not at all	₂ A little	₃ Somewhat	₄ Very much	₅ Extremely
124.	In general, how masculine do you behave in front of others?	₁ Not at all	₂ A little	₃ Somewhat	₄ Very much	₅ Extremely
125.	In general, how masculine do you appear to others?	₁ Not at all	₂ A little	₃ Somewhat	₄ Very much	₅ Extremely
The following questions are about your use of some different gender-affirming practices. We understand that not everyone does these practices; however, we appreciate any information you are able to share with us, whether you do these practices or not.						
139.	Do you use hormones for gender affirming care ("transitioning")?	₁ Yes, from a local private healthcare provider	₂ Yes, from a local public healthcare provider	₃ Yes, from another source	₀ No	
140.	Do you use any method of binding (binders, bandages, etc.)?				₁ Yes	₀ No
141.	Do you tuck (or use any method of hiding your penis)?				₁ Yes	₀ No

Section 1c: Sexuality and self

Complete this section if you do not identify as heterosexual or asexual. If you do identify as heterosexual or asexual, go to the next page.

Place an X in one box that best describes your answer to each question.

Please answer these questions based on YOUR OWN feelings about yourself.

126.	Sometimes I dislike myself for being a person who has (or wants) sex with people of the same sex.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
127.	I wish I was only sexually attracted to the opposite sex.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
128.	I am ashamed of myself for being sexually attracted to people of the same sex.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
129.	I feel that being attracted to people of the same sex is a personal weakness of mine.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
130.	If someone offered me the chance to be completely heterosexual, I would accept the offer.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
131.	Whenever I think about having sex with someone of the same sex, I feel bad about myself.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly

SWAZILAND—ENGLISH

Section 1d: Gender identity and self

Complete this section if you identify as transgender, genderqueer, and/or gender non-conforming. If you do not identify as transgender, genderqueer, and/or gender non-conforming, go to the next page.

Place an X in one box that best describes your answer to each question.

Please answer these questions based on YOUR OWN feelings about yourself.

132.	Sometimes I dislike myself for being transgender, genderqueer, and/or gender non-conforming.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
133.	Sometimes I wish I wasn't transgender, genderqueer, and/or gender non-conforming.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
134.	I think about the fact that I am transgender, genderqueer, and/or gender non-conforming when I interact with people.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
135.	I feel that being transgender, genderqueer, and/or gender non-conforming is a personal weakness of mine.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly
136.	If someone offered me the chance to be cisgender, I would accept the offer.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly

The following questions are about your access to gender-affirming treatments. We understand that not everyone chooses to use these treatments; however, we appreciate any information you are able to share with us about access, whether you use these treatments or not.

137.	Can you get hormones for transitioning from a local healthcare provider, if you need them?	1 Yes	0 No
138.	Can you get gender affirming surgery from a local healthcare provider, if you need it?	1 Yes	0 No

Section 1e: Being intersex and self

Complete this section if you are intersex. If you are not intersex, go to the next page.

Place an X in one box that best describes your answer to each question.

Please answer these questions based on YOUR OWN feelings about yourself.

142.	Sometimes I dislike myself for being intersex.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly	
143.	Sometimes I wish I wasn't intersex.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly	
144.	I think about the fact that I am intersex when I interact with people.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly	
145.	I feel that being intersex is a personal weakness of mine.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly	
146.	If someone offered me the chance to not have been born intersex, I would accept the offer.	1 Disagree strongly	2 Disagree	3 Agree	4 Agree strongly	
147.	How do you rate your healthcare providers' knowledge and skills on intersex healthcare?	4 Very good	3 Good	2 Poor	1 Very poor	
148.	Has healthcare staff ever put your body on display for others to look at?				1 Yes	0 No

Section 2a: Health service use

The following questions will ask about your health service use at community-based organisations/non-governmental organisations, public services, private services, and indigenous or traditional healers or providers.

201.	Do you have private medical aid or health insurance?	1 Yes	0 No
202.	<p>For which health services have you accessed community-based organisation, faith-based organisation, or non-governmental organisation healthcare in the last 12 months?</p> <p>TICK ALL THAT APPLY</p> <p><i>(If you do not use <u>community-based organisation or non-governmental organisation healthcare</u>, tick "None" at the bottom)</i></p>	<input type="checkbox"/> 1 Regular check-ups when I am feeling well <input type="checkbox"/> 2 Check-ups when I am feeling sick <input type="checkbox"/> 3 Emergency care <input type="checkbox"/> 4 Care after a sexual assault <input type="checkbox"/> 5 Care after a physical assault <input type="checkbox"/> 6 Testing for HIV <input type="checkbox"/> 7 HIV care and treatment <input type="checkbox"/> 8 Testing, care, or treatment for other sexually transmitted infections (STIs) (not HIV) <input type="checkbox"/> 15 Counselling or psychosocial support <input type="checkbox"/> 16 Care for mental health conditions <input type="checkbox"/> 10 Barrier methods (condoms, dental dams or finger condoms) <input type="checkbox"/> 11 Contraception (injection, pill, IUD/loop, implant) <input type="checkbox"/> 12 Gender affirming treatment (hormones, surgery) <input type="checkbox"/> 13 Other, specify: _____ <input type="checkbox"/> 14 None	
203.	<p>For which health services have you accessed public health care (clinic/hospital) in the last 12 months?</p> <p>TICK ALL THAT APPLY</p> <p><i>(If you do not use <u>public healthcare</u>, tick "None" at the bottom)</i></p>	<input type="checkbox"/> 1 Regular check-ups when I am feeling well <input type="checkbox"/> 2 Check-ups when I am feeling sick <input type="checkbox"/> 3 Emergency care <input type="checkbox"/> 4 Care after a sexual assault <input type="checkbox"/> 5 Care after a physical assault <input type="checkbox"/> 6 Testing for HIV <input type="checkbox"/> 7 HIV care and treatment <input type="checkbox"/> 8 Testing, care, or treatment for other sexually transmitted infections (STIs) (not HIV) <input type="checkbox"/> 15 Counselling or psychosocial support <input type="checkbox"/> 16 Care for mental health conditions <input type="checkbox"/> 10 Barrier methods (condoms, dental dams or finger condoms) <input type="checkbox"/> 11 Contraception (injection, pill, IUD/loop, implant) <input type="checkbox"/> 12 Gender affirming treatment (hormones, surgery) <input type="checkbox"/> 13 Other, specify: _____ <input type="checkbox"/> 14 None	

<p>204.</p>	<p>For which health services have you accessed private health care (clinic/hospital) in the last 12 months?</p> <p>TICK ALL THAT APPLY</p> <p><i>(If you do not use private healthcare, tick "None" at the bottom)</i></p>	<p><input type="checkbox"/> 1 Regular check-ups when I am feeling well</p> <p><input type="checkbox"/> 2 Check-ups when I am feeling sick</p> <p><input type="checkbox"/> 3 Emergency care</p> <p><input type="checkbox"/> 4 Care after a sexual assault</p> <p><input type="checkbox"/> 5 Care after a physical assault</p> <p><input type="checkbox"/> 6 Testing for HIV</p> <p><input type="checkbox"/> 7 HIV care and treatment</p> <p><input type="checkbox"/> 8 Testing, care, or treatment for other sexually transmitted infections (STIs) (not HIV)</p> <p><input type="checkbox"/> 15 Counselling or psychosocial support</p> <p><input type="checkbox"/> 16 Care for mental health conditions</p> <p><input type="checkbox"/> 10 Barrier methods (condoms, dental dams or finger condoms)</p> <p><input type="checkbox"/> 11 Contraception (injection, pill, IUD/loop, implant)</p> <p><input type="checkbox"/> 12 Gender affirming treatment (hormones, surgery)</p> <p><input type="checkbox"/> 13 Other, specify: _____</p> <p><input type="checkbox"/> 14 None</p>
<p>205.</p>	<p>For which health services have you accessed indigenous or traditional healthcare or faith healing in the last 12 months?</p> <p>TICK ALL THAT APPLY</p> <p><i>(If you do not use indigenous or traditional healthcare or faith healing, tick "None" at the bottom)</i></p>	<p><input type="checkbox"/> 1 Regular check-ups when I am feeling well</p> <p><input type="checkbox"/> 2 Check-ups when I am feeling sick</p> <p><input type="checkbox"/> 3 Emergency care</p> <p><input type="checkbox"/> 4 Care after a sexual assault</p> <p><input type="checkbox"/> 5 Care after a physical assault</p> <p><input type="checkbox"/> 6 Testing for HIV</p> <p><input type="checkbox"/> 7 HIV care and treatment</p> <p><input type="checkbox"/> 8 Testing, care, or treatment for other sexually transmitted infections (STIs) (not HIV)</p> <p><input type="checkbox"/> 15 Counselling or psychosocial support</p> <p><input type="checkbox"/> 16 Care for mental health conditions</p> <p><input type="checkbox"/> 10 Barrier methods (condoms, dental dams or finger condoms)</p> <p><input type="checkbox"/> 11 Contraception (injection, pill, IUD/loop, implant)</p> <p><input type="checkbox"/> 12 Gender affirming treatment (hormones, surgery)</p> <p><input type="checkbox"/> 13 Other, specify: _____</p> <p><input type="checkbox"/> 14 None</p>

SWAZILAND—ENGLISH

Section 2b: Health service barriers

Place an X in one box that best describes your answer to each question.

206.	Have you ever disclosed being LGBTI to a healthcare staff member? (this includes doctors, nurses, counsellors, other people working at public, private, or traditional healthcare facilities)			1 Yes	0 No
207.	Has a healthcare staff member ever made assumptions about your sexual orientation and/or gender identity? (for example, assumed you are LGBTI based on how you dress)			1 Yes	0 No
208.	When seeking healthcare, how often do you think you have been treated disrespectfully by staff for being LGBTI? (this includes doctors, nurses, counsellors, other people working at public, private, or traditional healthcare facilities)	1 Never	2 Rarely	3 Sometimes	4 Often
209.	When seeking healthcare, how often do you think have you received poorer service than other people for being LGBTI?	1 Never	2 Rarely	3 Sometimes	4 Often
210.	How often have you been called names or insulted by healthcare staff for being LGBTI? (this includes doctors, nurses, counsellors, other people working at public, private, or traditional healthcare facilities)	1 Never	2 Rarely	3 Sometimes	4 Often
211.	How often do you think healthcare staff has denied you a service because of being LGBTI? (this includes doctors, nurses, counsellors, other people working at public, private, or traditional healthcare facilities)	1 Never	2 Rarely	3 Sometimes	4 Often
212.	How often has healthcare staff threatened to call the police because you were LGBTI? (this includes doctors, nurses, counsellors, other people working at public, private, or traditional healthcare facilities)	1 Never	2 Rarely	3 Sometimes	4 Often
213.	Have you ever not told a healthcare staff member about a health need you have which is related to the fact that you are LGBTI? (for example, anal warts, sexual health advice for lesbian couples, gender-affirming treatment)			1 Yes	0 No

Section 2c: Impact of previous experiences on health-seeking behaviour

Place an X in one box that best describes your answer to each question.

214.	Have you postponed or not tried to get needed healthcare <u>when you were sick or injured</u> because you could not afford it?			1 Yes	0 No
215.	Have you postponed or not tried to get <u>HIV testing</u> because you could not afford it?			1 Yes	0 No
216.	Have you postponed or not tried to get <u>STI testing or STI/HIV treatment</u> because you could not afford it?			1 Yes	0 No
217.	Have you postponed or not tried to get needed healthcare <u>when you were sick or injured</u> because of disrespect or discrimination based on being LGBTI from doctors or other healthcare providers?			1 Yes	0 No
218.	Have you postponed or not tried to get <u>HIV testing</u> because of disrespect or discrimination based on being LGBTI from doctors or other healthcare providers?			1 Yes	0 No
219.	Have you postponed or not tried to get <u>STI testing or STI/HIV treatment</u> because of disrespect or discrimination based on being LGBTI from doctors or other healthcare providers?			1 Yes	0 No
220.	Have you ever hidden, or tried to hide, that you are LGBTI from a healthcare provider for fear of discrimination?			1 Yes	0 No
221.	Are you aware of a healthcare professional ever sharing that you are LGBTI with others without your permission?			1 Yes	0 No

Section 3: Tobacco

3001.	Do you currently smoke tobacco every day, some days, or not at all?	² Every day (Go to 3004)	¹ Some days (Go to 3002)	⁰ Not at all (Go to 3003)
3002.	Have you smoked tobacco every day in the past?		¹ Yes (Go to 3004)	⁰ No (Go to 3004)
3003.	In the past, have you ever smoked tobacco?	² Yes, every day in the past (Go to next section)	¹ Yes, some days in the past (Go to next section)	⁰ No (Go to next section)
3004.	On average, how many cigarettes do you currently smoke each day when you smoke?	Write the number per day: _____ Note: 1 pack = 20 cigarettes		

Section 3a: Alcohol

Because alcohol use can affect your health and can interfere with certain medications and treatments, it is important that we ask some questions about your use of alcohol. Your answers will remain confidential so please be honest.

Place an X in one box that best describes your answer to each question.

301.	How often do you have a drink containing alcohol?	⁰ Never (Go to next section)	¹ Monthly or less	⁽²⁾ 2-4 times a month	⁽³⁾ 2-3 times a week	⁽⁴⁾ 4 or more times a week
302.	How many drinks containing alcohol do you have on a typical day when you are drinking?	⁽⁰⁾ 1 or 2	⁽¹⁾ 3 or 4	⁽²⁾ 5 or 6	⁽³⁾ 7, 8 or 9	⁽⁴⁾ 10 or more
303.	How often do you have six or more drinks on one occasion?	⁰ Never	¹ Less than monthly	² Monthly	³ Weekly	⁴ Daily or almost daily
304.	How often during the last year have you found that you were not able to stop drinking once you had started?	⁰ Never	¹ Less than monthly	² Monthly	³ Weekly	⁴ Daily or almost daily
305.	How often during the last year have you failed to do what was normally expected of you because of drinking?	⁰ Never	¹ Less than monthly	² Monthly	³ Weekly	⁴ Daily or almost daily
306.	How often during the last year have you needed a first drink in the morning to get yourself going after a heavy drinking session?	⁰ Never	¹ Less than monthly	² Monthly	³ Weekly	⁴ Daily or almost daily
307.	How often during the last year have you had a feeling of guilt or remorse after drinking?	⁰ Never	¹ Less than monthly	² Monthly	³ Weekly	⁴ Daily or almost daily
308.	How often during the last year have you been unable to remember what happened the night before because of your drinking?	⁰ Never	¹ Less than monthly	² Monthly	³ Weekly	⁴ Daily or almost daily
309.	Have you or someone else been injured because of your drinking?	⁰ No		² Yes, but not in the last year		⁴ Yes, during the last year
310.	Has a relative, friend, doctor, or other health care worker been concerned about your drinking or suggested you cut down?	⁰ No		² Yes, but not in the last year		⁴ Yes, during the last year

Section 3b: Drugs

Here are a few questions about drugs. Please answer as correctly and honestly as possible.

By drugs, we mean any of the following:

Cannabis: Marijuana, Hash, Hash oil, Dagga

Amphetamines: Methamphetamine, Phenmetraline, Khat, Betel nut, Ritaline, (Methylphenidate)

Cocaine: Crack, Freebase, Coca leaves

Opiates: Smoked heroin, Heroin, Opium

Hallucinogens: Ecstasy, LSD (Lisergic acid), Mescaline, Peyote, PCP (angel dust), (Phencyclidine), Psilocybin, DMT (Dimethyltryptamine)

Solvents/inhalants: Thinner, Trichlorethylene, Gasoline/petrol, Gas, Solution, Glue

GHB and others: GHB, Anabolic steroids, Laughing gas (Halothane), Amyl nitrate (Poppers), Anticholinergic compounds

Tik or rocks

Place an X in one box that best describes your answer to each question.

311.	How often do you use drugs other than alcohol? (see list of drugs above)	<input type="radio"/> Never (Go to next section)	<input type="radio"/> 1 Once a month or less often	<input type="radio"/> 2 2-4 times a month	<input type="radio"/> 3 2-3 times a week	<input type="radio"/> 4 4 times a week or more often
312.	Do you use more than one type of drug on the same occasion?	<input type="radio"/> Never	<input type="radio"/> 1 Once a month or less often	<input type="radio"/> 2 2-4 times a month	<input type="radio"/> 3 2-3 times a week	<input type="radio"/> 4 4 times a week or more often
313.	How many times do you take drugs on a typical day when you use drugs?	<input type="radio"/> 0	<input type="radio"/> 1 1-2	<input type="radio"/> 2 3-4	<input type="radio"/> 3 5-6	<input type="radio"/> 4 7 or more
314.	How often are you influenced heavily by drugs?	<input type="radio"/> Never	<input type="radio"/> 1 Less often than once a month	<input type="radio"/> 2 Every month	<input type="radio"/> 3 Every week	<input type="radio"/> 4 Daily or almost every day
315.	Over the past year, have you felt that your longing for drugs was so strong that you could not resist it?	<input type="radio"/> Never	<input type="radio"/> 1 Less often than once a month	<input type="radio"/> 2 Every month	<input type="radio"/> 3 Every week	<input type="radio"/> 4 Daily or almost every day
316.	Has it happened, over the past year that you have not been able to stop taking drugs once you started?	<input type="radio"/> Never	<input type="radio"/> 1 Less often than once a month	<input type="radio"/> 2 Every month	<input type="radio"/> 3 Every week	<input type="radio"/> 4 Daily or almost every day
317.	How often over the past year have you taken drugs and then neglected to do something you should have done?	<input type="radio"/> Never	<input type="radio"/> 1 Less often than once a month	<input type="radio"/> 2 Every month	<input type="radio"/> 3 Every week	<input type="radio"/> 4 Daily or almost every day
318.	How often over the past year have you needed to take a drug the morning after heavy drug use the day before?	<input type="radio"/> Never	<input type="radio"/> 1 Less often than once a month	<input type="radio"/> 2 Every month	<input type="radio"/> 3 Every week	<input type="radio"/> 4 Daily or almost every day
319.	How often over the past year have you had guilty feelings or a bad conscience because you used drugs?	<input type="radio"/> Never	<input type="radio"/> 1 Less often than once a month	<input type="radio"/> 2 Every month	<input type="radio"/> 3 Every week	<input type="radio"/> 4 Daily or almost every day
320.	Have you or anyone else been hurt (mentally or physically) because you used drugs?	<input type="radio"/> No		<input type="radio"/> 2 Yes, but not over the past year		<input type="radio"/> 4 Yes, over the past year
321.	Has a relative or a friend, a doctor, or a nurse, or anyone else, been worried about your drug use?	<input type="radio"/> No		<input type="radio"/> 2 Yes, but not over the past year		<input type="radio"/> 4 Yes, over the past year

Section 3c

Over the last 2 weeks, how often have you been bothered by the following problems?

322.	Feeling nervous, anxious, or on edge	⁰ Not at all (0-1 days)	¹ Several days (2-6 days)	² Over half the days (7-10 days)	³ Nearly every day (11-14 days)	
323.	Not being able to stop or control worrying	⁰ Not at all (0-1 days)	¹ Several days (2-6 days)	² Over half the days (7-10 days)	³ Nearly every day (11-14 days)	
324.	Worrying too much about different things	⁰ Not at all (0-1 days)	¹ Several days (2-6 days)	² Over half the days (7-10 days)	³ Nearly every day (11-14 days)	
325.	Trouble relaxing	⁰ Not at all (0-1 days)	¹ Several days (2-6 days)	² Over half the days (7-10 days)	³ Nearly every day (11-14 days)	
326.	Being so restless that it is hard to sit still	⁰ Not at all (0-1 days)	¹ Several days (2-6 days)	² Over half the days (7-10 days)	³ Nearly every day (11-14 days)	
327.	Becoming easily annoyed or irritable	⁰ Not at all (0-1 days)	¹ Several days (2-6 days)	² Over half the days (7-10 days)	³ Nearly every day (11-14 days)	
328.	Feeling afraid as if something awful might happen	⁰ Not at all (0-1 days)	¹ Several days (2-6 days)	² Over half the days (7-10 days)	³ Nearly every day (11-14 days)	
329.	If you checked off any problems, how difficult have these made it for you to do your work, take care of things at home, or get along with other people?	⁰ Not difficult at all	¹ Somewhat difficult	² Very difficult	³ Extremely difficult	
330.	Has a healthcare provider ever told you that you have clinical anxiety?				¹ Yes	⁰ No (Go to next section)
330a.	If yes, are you current being treated for clinical anxiety (e.g. medication, therapy)?				¹ Yes	⁰ No

SWAZILAND—ENGLISH

Section 3d

Below is a list of some of the ways you may have felt or behaved. Please indicate how often you have felt this way **during the past week**.

331.	I was bothered by things that usually don't bother me.	⁰ Rarely or none of the time (less than 1 day)	¹ Some or a little of the time (1-2 days)	² Occasionally or a moderate amount of time (3-4 days)	³ All of the time (5-7 days)
332.	I had trouble keeping my mind on what I was doing.	⁰ Rarely or none of the time (less than 1 day)	¹ Some or a little of the time (1-2 days)	² Occasionally or a moderate amount of time (3-4 days)	³ All of the time (5-7 days)
333.	I felt depressed.	⁰ Rarely or none of the time (less than 1 day)	¹ Some or a little of the time (1-2 days)	² Occasionally or a moderate amount of time (3-4 days)	³ All of the time (5-7 days)
334.	I felt that everything I did was an effort.	⁰ Rarely or none of the time (less than 1 day)	¹ Some or a little of the time (1-2 days)	² Occasionally or a moderate amount of time (3-4 days)	³ All of the time (5-7 days)
335.	I felt hopeful about the future.	³ Rarely or none of the time (less than 1 day)	² Some or a little of the time (1-2 days)	¹ Occasionally or a moderate amount of time (3-4 days)	⁰ All of the time (5-7 days)
336.	I felt fearful.	⁰ Rarely or none of the time (less than 1 day)	¹ Some or a little of the time (1-2 days)	² Occasionally or a moderate amount of time (3-4 days)	³ All of the time (5-7 days)
337.	My sleep was restless.	⁰ Rarely or none of the time (less than 1 day)	¹ Some or a little of the time (1-2 days)	² Occasionally or a moderate amount of time (3-4 days)	³ All of the time (5-7 days)
338.	I was happy.	³ Rarely or none of the time (less than 1 day)	² Some or a little of the time (1-2 days)	¹ Occasionally or a moderate amount of time (3-4 days)	⁰ All of the time (5-7 days)
339.	I felt lonely.	⁰ Rarely or none of the time (less than 1 day)	¹ Some or a little of the time (1-2 days)	² Occasionally or a moderate amount of time (3-4 days)	³ All of the time (5-7 days)
340.	I could not "get going."	⁰ Rarely or none of the time (less than 1 day)	¹ Some or a little of the time (1-2 days)	² Occasionally or a moderate amount of time (3-4 days)	³ All of the time (5-7 days)
341.	Has a healthcare provider ever told you that you have clinical depression?			¹ Yes	⁰ No (Go to 342)
341a.	If yes, are you current being treated for clinical depression (e.g. medication, therapy)?			¹ Yes	⁰ No

Section 3e

342.	Has there ever been a period of time when you thought about committing suicide?	a. In your lifetime?	¹ Yes	⁰ No
		b. In the last 12 months?	¹ Yes	⁰ No
343.	Did you ever try to end your own life, whether or not you had thought about it ahead?	a. In your lifetime?	¹ Yes	⁰ No
		b. In the last 12 months?	¹ Yes	⁰ No

Section 3f: Social support

<p>347.</p>	<p>Who do you go to when you need someone to talk to about problems in your life?</p> <p><u>TICK ALL THAT APPLY</u></p>	<p><input type="checkbox"/> 1 Current partner(s) (at least one)</p> <p><input type="checkbox"/> 2 Family (at least one member)</p> <p><input type="checkbox"/> 3 Friends (at least one)</p> <p><input type="checkbox"/> 4 People I live with (at least one)</p> <p><input type="checkbox"/> 5 Healthcare providers (at least one)</p> <p><input type="checkbox"/> 6 People I work with (at least one)</p> <p><input type="checkbox"/> 7 People living nearby me (at least one)</p> <p><input type="checkbox"/> 8 LGBTI organisations</p> <p><input type="checkbox"/> 9 No one</p>
<p>348.</p>	<p>Who in your life knows that you are LGBTI?</p> <p><u>TICK ALL THAT APPLY</u></p>	<p><input type="checkbox"/> 1 Current partner(s) (at least one)</p> <p><input type="checkbox"/> 2 Family (at least one member)</p> <p><input type="checkbox"/> 3 Friends (at least one)</p> <p><input type="checkbox"/> 4 People I live with (at least one)</p> <p><input type="checkbox"/> 5 Healthcare providers (at least one)</p> <p><input type="checkbox"/> 6 People I work with (at least one)</p> <p><input type="checkbox"/> 7 People living nearby me (at least one)</p> <p><input type="checkbox"/> 8 LGBTI organisations</p> <p><input type="checkbox"/> 9 No one</p>
<p>349.</p>	<p>Of those, who have <u>you</u> told yourself about being LGBTI?</p> <p><u>TICK ALL THAT APPLY</u></p>	<p><input type="checkbox"/> 1 Current partner(s) (at least one)</p> <p><input type="checkbox"/> 2 Family (at least one member)</p> <p><input type="checkbox"/> 3 Friends (at least one)</p> <p><input type="checkbox"/> 4 People I live with (at least one)</p> <p><input type="checkbox"/> 5 Healthcare providers (at least one)</p> <p><input type="checkbox"/> 6 People I work with (at least one)</p> <p><input type="checkbox"/> 7 People living nearby me (at least one)</p> <p><input type="checkbox"/> 8 LGBTI organisations</p> <p><input type="checkbox"/> 9 No one</p>

SWAZILAND—ENGLISH

Section 4 Experience of violence

This is the last section of the questionnaire. The following questions ask about your experiences with violence.

401.	Are you aware of anyone ever revealing that you are LGBTI to others without your permission?		1 Yes	0 No	
402.	Has anyone ever threatened to reveal that you are LGBTI to others without your permission?		1 Yes	0 No	
403.	Has anyone ever insulted or verbally harassed you because of being LGBTI?	a. In your life time?	1 Yes	0 No	
		b. In the last 12 months?	1 Yes	0 No	
404.	Has an intimate partner (past or current) ever threatened to reveal that you are LGBTI to others without your permission?		1 Yes	0 No	
405.	Has an intimate partner (past or current) ever made you feel worthless because of being LGBTI?		1 Yes	0 No	
406.	Has an intimate partner (past or current) ever made you feel ashamed because of being LGBTI?		1 Yes	0 No	
407.	Have you ever been coerced, pressured or forced into marriage?		1 Yes	0 No	
408.	Have you ever been sexually assaulted	By an intimate partner of the same sex as you?	a. In your life time?	1 Yes	0 No
			b. In the last 12 months?	1 Yes	0 No
		By an intimate partner of a different sex than you?	c. In your life time?	1 Yes	0 No
			d. In the last 12 months?	1 Yes	0 No
		By someone you know (not an intimate partner but a neighbour, friend, family member, etc.)	e. In your life time?	1 Yes	0 No
			f. In the last 12 months?	1 Yes	0 No
		By a stranger	g. In your life time?	1 Yes	0 No
			h. In the last 12 months?	1 Yes	0 No
		By someone you live with? (an intimate partner or other person)	i. In your life time?	1 Yes	0 No
			j. In the last 12 months?	1 Yes	0 No
409.	Have you ever been physically assaulted	By an intimate partner of the same sex as you?	a. In your life time?	1 Yes	0 No
			b. In the last 12 months?	1 Yes	0 No
		By an intimate partner of a different sex than you?	c. In your life time?	1 Yes	0 No
			d. In the last 12 months?	1 Yes	0 No
		By someone you know (not an intimate partner but a neighbour, friend, family member, etc.)	e. In your life time?	1 Yes	0 No
			f. In the last 12 months?	1 Yes	0 No
		By a stranger	g. In your life time?	1 Yes	0 No
			h. In the last 12 months?	1 Yes	0 No
		By someone you live with? (an intimate partner or other person)	i. In your life time?	1 Yes	0 No
			j. In the last 12 months?	1 Yes	0 No

SWAZILAND—ENGLISH

If you answered yes to sexual or physical assault in your life time, please complete these questions:

We know that our sexual orientation and gender identity is not always easily separated. However, please choose the best response to these last questions.			
413.	Do you think any of these incidents (sexual or physical assault) were motivated by your sexual orientation?	<input type="radio"/> Yes	<input type="radio"/> No
414.	Do you think any of these incidents (sexual or physical assault) were motivated by your gender identity?	<input type="radio"/> Yes	<input type="radio"/> No
415.	Do you think any of these incidents (sexual or physical assault) were motivated by your body being intersex or not typically female/typically male?	<input type="radio"/> Yes	<input type="radio"/> No
416.	Did any of these incidents result in flashbacks, nightmares, or reliving the event?	<input type="radio"/> Yes	<input type="radio"/> No
417.	Have you avoided situations or people who remind you of the incident(s)?	<input type="radio"/> Yes	<input type="radio"/> No
418.	Following the incident(s), have you felt jumpy, irritable, or restless?	<input type="radio"/> Yes	<input type="radio"/> No

If you answered yes to sexual or physical assault in the last 12 months, please complete these questions:

410.	If you have experienced physical or sexual assault in the last 12 months, have you sought medical care for it?	<input type="radio"/> Yes	<input type="radio"/> No			
411.	If you have experienced physical or sexual assault in the last 12 months, have you reported it to the police ?	<input type="radio"/> Yes	<input type="radio"/> No			
412.	When seeking help for physical or sexual assault, how often do you think you have been treated with less courtesy than other people by police or healthcare staff for being LGBTI?	<input type="radio"/> Never	<input type="radio"/> Rarely	<input type="radio"/> Sometimes	<input type="radio"/> Often	<input type="radio"/> I have not sought help for physical or sexual assault

Thank you for your time in completing this survey! Please take a moment to check you have completed all of the questions.

Return this survey to the person who gave it to you when you are finished.

Thank you for telling us about your experiences of mental health, drug/alcohol use, and violence. If you would like to talk to someone about these things, please contact one of the below organisations:

Organisation	Contact details
Lifeline/Childline	Tel: (+27 11) 715-2000 Dicon number (switchboard cell) : 082-231-0805 Services: LifeLine offers individual counselling both face-to-face and by a 24-hour telephone counselling.
The Manzini Psychiatric Centre	Tel:+2682 505 5170 P.O Box 424, Manzini Services: drugs/alcohol counselling
Swagaa	Tel: +2682 5052 2899 (Helpline 951) P.O Box 560 Matsapha Services: violence counselling

SWAZILAND—ENGLISH

Kamhlaba	+2682 505 3586 ext.2023 P.O Box 1051 Manzini Services: psychosocial support
Swaziland AIDS Support organisation	+2682 2079943 P.O Box 838 Nhlngano Services: Sexual health counselling
SWAPOL Swaziland for Positive Living	Tel:+268 505 7088 or +268 505 6172 Nkoseluhlaza street, Mcozini building first floor, Office no. 9 Manzini Services: LGBTI support
The Rock of Hope	Tel: +2682 505 3586 ext. 2025 1228 Mbabha Street Services: LGBTI support
House of Our Pride (HOOP)	Services: LGBTI support

For research staff use only:

I, the **fieldworker**, have reviewed this questionnaire for completeness and accuracy.

Fieldworker signature: _____ Date: _____

I, the **research coordinator (or designee)**, have reviewed this questionnaire for completeness and accuracy.

Coordinator/designee signature: _____ Date: _____

I, the **GHJRU research staff member**, have reviewed this questionnaire for completeness and accuracy.

GHJRU signature: _____ Date: _____

I, the **data enterer**, have completed data entry of this questionnaire and assigned a unique identifier.

Data enterer signature: _____ Date: _____

Notes

